

Donkey Anti-Rabbit IgG H&L (DyLight® 488) preadsorbed ab96919

★★★★★ [1 Abreviews](#) [20 References](#) [画像数 4](#)

製品の概要

製品名	Donkey Anti-Rabbit IgG H&L (DyLight® 488) preadsorbed
由来種	Donkey
ターゲット生物種	Rabbit
特異性	By immunoelectrophoresis and ELISA this antibody reacts specifically with Rabbit IgG and with light chains common to other Rabbit immunoglobulins. No antibody was detected against non immunoglobulin serum proteins. Reduced cross-reactivity to bovine, chicken, goat, horse, human, mouse, pig and rat was detected.
アプリケーション	適用あり: WB, IHC-P, ICC/IF, Flow Cyt
吸着処理血清	Chicken, Cow, Goat, Horse, Human, Mouse, Pig, Rat more details
標識	DyLight® 488. Ex: 493nm, Em: 518nm

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	pH: 6.8 Preservative: 0.09% Sodium azide Constituents: 0.2% BSA, PBS
精製度	Immunogen affinity purified
特記事項(精製)	Antiserum was cross adsorbed using bovine, chicken, goat, human, mouse, pig and rat immunosorbents to remove cross reactive antibodies. This antibody was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 488.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab96919の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		1/1000 - 1/20000. Predicted molecular weight: 36 kDa.
IHC-P		1/50 - 1/500.
ICC/IF	★★★★★ (1)	1/50 - 1/500.
Flow Cyt		1/50 - 1/200.

画像




rabbit anti-actin antibody at 1 µg/ml + HeLa whole cell lysate at 30 µg

Secondary

Donkey Anti-Rabbit IgG H&L (DyLight® 488) preadsorbed (ab96919) at 1/1000 dilution (Donkey anti-rabbit IgG heavy and light chain cross-adsorbed antibody)


Predicted band size: 36 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Donkey Anti-Rabbit IgG H&L (DyLight® 488) preadsorbed (ab96919)

This image is courtesy of an abreview submitted by Carl Hobbs, Kings's College London, United Kingdom.


Glial fibrillary acidic protein (GFAP) staining in mouse Formalin Fixed Paraffin Embedded sections. For antigen retrieval HIER in citrate buffer was performed. Donkey polyclonal Secondary Antibody to Rabbit IgG - H&L (DyLight® 488), pre-adsorbed (ab96919) was used as a secondary antibody at a dilution of 1/50.



Immunohistochemistry (Frozen sections) - Donkey Anti-Rabbit IgG H&L (DyLight® 488) preadsorbed (ab96919)

E14 mouse brain sample was stained with primary antibody (**ab66155**). The secondary antibody used was ab96919 (1:200 dilution, incubate for 1 hour at room temperature). The blue background was from DAPI.

This image is courtesy of an anonymous Abreview



Donkey Anti-Rabbit IgG H&L (DyLight® 488) preadsorbed (ab96919)

Emission spectra of DyLight® fluorochromes available in our catalog.

Line colors represent the approximate visible colors of the wavelength maxima.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors