abcam

Product datasheet

Goat Anti-Mouse IgG H&L (DyLight® 594) preadsorbed ab96881

23 References 画像数 2

製品の概要

製品名 Goat Anti-Mouse IgG H&L (DyLight® 594) preadsorbed

由来種Goatターゲット生物種Mouse

特異性 By immunoelectrophoresis and ELISA this antibody reacts specifically with Mouse IgG and with

light chains common to other Mouse immunoglobulins. No antibody was detected against non immunoglobulin serum proteins. Reduced cross-reactivity to bovine, chicken, goat, horse, human,

pig, rabbit and rat lgG was detected.

アプリケーション 適用あり: ICC/IF, IHC-P, Flow Cyt

吸着処理血清

Chicken, Cow, Goat, Horse, Human, Pig, Rabbit, Rat more details

免疫原 Other Immunogen Type corresponding to Mouse IgG.

標識 DyLight® 594. Ex: 593nm, Em: 618nm

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C.

バッファー pH: 6.8

Preservative: 0.09% Sodium azide Constituents: 0.2% BSA, PBS

精製度 Immunogen affinity purified

特記事項(精製) Antiserum was cross adsorbed using bovine, chicken, horse, human, pig, rabbit and rat

immunosorbents to remove cross reactive antibodies. This antibody was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 594.

ポリ/モノ ポリクローナル

アイソタイプ IgG


アプリケーション

1

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab96881の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
ICC/IF		1/50 - 1/500.
IHC-P		1/50 - 1/500.
Flow Cyt		1/50 - 1/200.

画像



Emission spectra of DyLight® fluorochromes available in our catalog.

Line colors represent the approximate visible colors of the wavelength maxima.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Goat Anti-Mouse IgG H&L (DyLight® 594) preadsorbed (ab96881)

ab96881 was used at dilution 1/100 for IHC analysis of FFPE human kidney. Primary Antibody: mouse anti-p53 (clone DO-1) used at a dilution of 1/100. Secondary Antibody: Red-fluorescent goat anti-mouse lgG-heavy and light chain cross-adsorbed Antibody DyLight 594 Conjugated ab96881 (2 different lots) used at a dilution of 1/100 (5 μ g/mL). Counterstain: DAPI (blue)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors