

Goat Anti-Mouse IgG H&L (FITC) ab6785

★★★★★ [1 Abreviews](#)   [510 References](#)   [画像数 4](#)

製品の概要

製品名	Goat Anti-Mouse IgG H&L (FITC)
由来種	Goat
ターゲット生物種	Mouse
アプリケーション	<b>適用あり:</b> IHC (PFA fixed), Immunomicroscopy, Flow Cyt, IHC-P, IHC-Fr, ICC/IF, ELISA, WB
免疫原	mouse whole molecule
標識	FITC. Ex: 493nm, Em: 528nm

製品の特性


製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	pH: 6.50 Preservative: 0.01% Sodium azide Constituents: 0.42% Tripotassium orthophosphate, 0.87% Sodium chloride, 1% BSA
精製度	Affinity purified
特記事項 (精製)	This product was prepared from monospecific antiserum by immunoaffinity chromatography using mouse IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities.
特記事項 (標識)	Diaminotriazinylaminofluorescein (DTAF) (Molecular Weight 530 daltons) Absorption Wavelength: 495 nm Emission Wavelength: 528 nm Fluorochrome/Protein Ratio: 3.3 moles FITC per mole of Goat IgG
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

**The Abpromise guarantee**      **Abpromise保証は、**[次のテスト済みアプリケーション](#)におけるab6785の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC (PFA fixed)		1/1000.
Immunomicroscopy		Use at an assay dependent concentration.
Flow Cyt		1/500 - 1/2500. PubMed: 17989178
IHC-P		Use at an assay dependent concentration.
IHC-Fr		Use at an assay dependent concentration. PubMed: 15716365 For PFA fixed tissue see PMID 15716365.
ICC/IF	★★★★★ (1)	1/1000 - 1/5000. (PMID16006199)
ELISA		1/10000 - 1/50000.
WB		1/10000. PubMed: 17217624


## 画像



Immunocytochemistry/ Immunofluorescence - Goat  
Anti-Mouse IgG H&L (FITC) (ab6785)


This image is courtesy of an Abreview submitted by J Chai.

**ab8978** staining vimentin in human colon fibroblasts by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with methanol, permeabilized with 0.1% Triton X-100 and blocked with Dako serum free protein blocker for 20 minutes at 28°C. Samples were incubated with primary antibody (1/100 in Dako antibody diluent) for 2 hours at 28°C. A FITC-conjugated goat anti-mouse IgG H&L (ab6785) (1/800) was used as the secondary antibody.




Flow Cytometry - Goat Anti-Mouse IgG H&L (FITC)  
(ab6785)

Ab6785 was used at dilution 1/100 with the primary antibody [ab52455](#) in Flow Cyt. See the review on [ab52455](#).



Immunocytochemistry/ Immunofluorescence - Goat  
Anti-Mouse IgG H&L (FITC) (ab6785)

Ab6785 was used at dilution 1/2000 with the primary antibody [ab1220](#) in ICC/IF. See the review on [ab1220](#).



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Goat Anti-Mouse IgG H&L (FITC) (ab6785)

Ab6785 was used at dilution 1/300 with the primary antibody **ab6002** in IHC-P. See the review on **ab6002**.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors