

### Goat Anti-Rabbit IgG H&L (Cy2 ®) preadsorbed ab6940

★★★★★ [1 Abreviews](#) [39 References](#) [画像数 3](#)

#### 製品の概要

製品名	Goat Anti-Rabbit IgG H&L (Cy2 ®) preadsorbed
由来種	Goat
ターゲット生物種	Rabbit
アプリケーション	<b>適用あり:</b> WB, ICC/IF, IHC-P, IHC-Fr, ICC, Flow Cyt, ELISA
吸着処理血清	Chicken, Cow, Goat, Guinea pig, Hamster, Horse, Human, Mouse, Rat, Sheep <a href="#">more details</a>
免疫原	Full length native Rabbit IgG (purified).
標識	Cy2 ®. Ex: 489nm, Em: 505nm

#### 製品の特性


製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	pH: 6.50 Preservative: 0.01% Sodium azide Constituents: 0.42% Tripotassium orthophosphate, 0.87% Sodium chloride, 1% BSA
精製度	Immunogen affinity purified
特記事項 (精製)	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads followed by solid phase adsorption(s) to remove any unwanted reactivities and extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum, Rabbit IgG and Rabbit Serum.
特記事項 (標識)	Cy2 (Cyanine 2-OSu) (Molecular Weight 897 daltons) Absorption Wavelength: 489 nm Emission Wavelength: 505 nm Fluorochrome/Protein Ratio: 2.0 moles Cy2 per mole of Goat IgG
ポリ/モノ	ポリクローナル
アイソタイプ	IgG
特記事項	Cy™ and CyDye™ are registered trademarks of Cytiva.  This secondary antibody is specifically designed for the detection of multiple primary antibodies (polyclonal or monoclonal) of different host species in experiments where cells are simultaneously labeled without unwanted cross reaction.

## アプリケーション

**The Abpromise guarantee** Abpromise保証は、次のテスト済みアプリケーションにおけるab6940の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent dilution.
ICC/IF		1/1000 - 1/5000.
IHC-P		Use at an assay dependent concentration.
IHC-Fr		Use at an assay dependent concentration.
ICC	★★★★★ (1)	Use at an assay dependent concentration.
Flow Cyt		Use at an assay dependent dilution.
ELISA		1/10000 - 1/50000.

## 画像



Goat Anti-Rabbit IgG H&L (Cy2®) preadsorbed (ab6940) at  
1/1000 dilution + Rabbit IgG at 0.05 µg

Incubated with ab6940 for 1 hour at room temperature.



**All lanes :** Goat Anti-Rabbit IgG H&L (Cy2 ®) preadsorbed (ab6940) at 1/1000 dilution

**Lane 1 :** Rabbit IgG whole molecule

**Lane 2 :** Rabbit IgG F(ab) fragment

**Lane 3 :** Rabbit IgG F(c) fragment

**Lane 4 :** Rabbit IgM whole molecule

**Lane 5 :** Normal rabbit serum

Lysates/proteins at 0.05 µg per lane.

### Secondary


**All lanes :** Peroxidase-conjugated donkey anti-goat IgG at 1/40000 dilution

Performed under reducing conditions.

**Observed band size:** 23,25,50,70 kDa

Incubated with the primary antibody for 1 hour at room temperature.

Incubated with the secondary antibody for 30 minutes at room temperature.



ab6940 was used at dilution 1/10000 with the primary antibody **ab47840** in WB. See the review on **ab47840**.

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

#### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors