

Mouse spleen tissue lysate - total protein ab7937

3 References

製品の概要

製品名 Mouse spleen tissue lysate - total protein

特記事項 Source: White laboratory mice

Mouse spleen tissue lysate was prepared by homogenization in modified RIPA buffer (150mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM phenylmethylsulfonyl fluoride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecylsulfate, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin). Tissue and cell debris were removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The tissue lysate was boiled for 5 min in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol.

Abcam has not and does not intend to apply for the REACH Authorisation of customers' uses of products that contain European Authorisation list (Annex XIV) substances. It is the responsibility of our customers to check the necessity of application of REACH Authorisation, and any other relevant authorisations, for their intended uses.

アプリケーション **適用あり:** WB

製品の特性

Mycoplasma free Yes

製品の状態 Liquid

保存方法 Shipped at 4°C. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle.

バッファー pH: 7.2
Constituents: 60.05% Water, 12.5% Glycerol (glycerin, glycerine), 9% Tris HCl, 7.7% DTT, 4.4% Sodium chloride, 1% Triton-X-100, 1% Sodium deoxycholate, 1.1% Sodium lauryl sulfate, 0.15% EDTA disodium salt, 0.5% Aprotinin, 0.5% Leupeptin hemisulfate, 0.09% PMSF, 0.01% Bromophenol blue

ライゼート備考 Mouse spleen tissue lysate was prepared by homogenization in modified RIPA buffer (150mM sodium chloride, 50 mM Tris-HCl, pH 7.4, 1 mM ethylenediaminetetraacetic acid, 1 mM phenylmethylsulfonyl fluoride, 1% Triton X-100, 1% sodium deoxycholic acid, 0.1% sodium dodecylsulfate, 5 µg/ml of aprotinin, 5 µg/ml of leupeptin). Tissue and cell debris were removed by centrifugation. Protein concentration was determined with Bio-Rad protein assay. The tissue lysate was boiled for 5 min in 1 x SDS sample buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1%

sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol.

アプリケーション

The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab7937の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration. Mouse spleen tissue lysate is ready to load on SDS-PAGE for Western blotting, 20 µg per lane is recommended for mini gel.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors