

Product datasheet

(-)-Cannabidiol ab120448


画像数 1

製品の概要

| | |
|---------|---|
| 製品名 | (-)-Cannabidiol |
| 製品の詳細 | Common constituent of Cannabis |
| 生理活性の詳細 | Common constituent of Cannabis with much lower affinity at CB ₁ and CB ₂ receptors (K _i values are 4350 and 2860 nM, respectively) than Delta-9 THC. Elicits anticonvulsant, anticancer, anti-inflammatory and neuroprotective effects. Protects retinal neurons in diabetic models. |
| 精製度 | > 98% |

製品の特性

| | |
|-----|---|
| 体系名 | 2-[(1 <i>R</i> ,6 <i>R</i>)-3-Methyl-6-(1-methylethenyl)-2-cyclohexen-1-yl]-5-pentyl-1,3-benzenediol |
| 分子量 | 314.47 |
| 構造式 | |



| | |
|--------------|--|
| 分子式 | C ₂₁ H ₃₀ O ₂ |
| CAS 番号 | 13956-29-1 |
| PubChem 登録番号 | 644019 |
| 保存方法 | Store at -20°C. Store under desiccating conditions. The product can be stored for up to 12 months. |
| 溶解性 | Soluble in DMSO to 100 mM and in ethanol to 100 mM |
| 使用に関する注意 | Wherever possible, you should prepare and use solutions on the same day. However, if you need to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and prior to opening the vial we recommend that you allow your product to equilibrate to room temperature for at least 1 hour. |

Need more advice on solubility, usage and handling? Please visit our [frequently asked questions \(FAQ\) page](#) for more details.

SMILES 線形表記

CC(=C)[C@@H]2CCC(C)=C[C@H]2c1c(O)cc(CCCCC)cc1O

由来

Synthetic


アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab120448** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| アプリケーション | Abreviews | 特記事項 |
|--------------------|-----------|--|
| Functional Studies | | Use at an assay dependent concentration. |

画像



[ab109225](#) staining Nox4 in HeLa cells treated with (-)-cannabidiol ([ab120448](#)), by ICC/IF. Increase in Nox4 expression correlates with increased concentration of (-)-cannabidiol, as described in literature.

The cells were incubated at 37°C for 6h in media containing different concentrations of [ab120847](#) ((-)-cannabidiol) in DMSO, fixed with 4% formaldehyde for 10 minutes at room temperature and blocked with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% tween for 2h at room temperature. Staining of the treated cells with [ab109225](#) (5 μg/ml) was performed overnight at 4°C in PBS containing 1% BSA and 0.1% tween. A DyLight 488 goat anti-rabbit polyclonal antibody ([ab96899](#)) at 1/250 dilution was used as the secondary antibody.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE, NOT FOR USE IN HUMANS"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team