

Product datasheet

Anti-ABCB9 antibody - Carboxyterminal end ab60036

画像数 1

製品の概要

製品名	Anti-ABCB9 antibody - Carboxyterminal end
製品の詳細	Rabbit polyclonal to ABCB9 - Carboxyterminal end
由来種	Rabbit
アプリケーション	適用あり: ELISA, IHC-P
種交差性	交差種: Human
免疫原	Synthetic peptide (10-30 aa in length) at the C-terminal of last 50 aa of human ABCB9 conjugated to KLH
ポジティブ・コントロール	Human hepatocarcinoma tissue.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.09% Sodium Azide Constituents: PBS
精製度	Protein G purified
特記事項 (精製)	ab60036 is purified through a protein G column and eluted out with both high and low pH buffers and neutralized immediately after elution then followed by dialysis against PBS.
ポリモノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab60036** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


アプリケーション	Abreviews	特記事項
ELISA		1/1000.

アプリケーション	Abreviews	特記事項
IHC-P		1/50 - 1/100.

ターゲット情報

関連性	<p>The membrane associated protein encoded by this gene is a member of the superfamily of ATP binding cassette (ABC) transporters. ABC proteins transport various molecules across extra and intra cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the MDR/TAP subfamily. Members of the MDR/TAP subfamily are involved in multidrug resistance as well as antigen presentation. The function of this half transporter has not yet been determined; however, this protein may play a role in lysosomes. Alternative splicing of this gene results in three known products which are likely to have different substrate specifications.</p>
細胞内局在	<p>Integral membrane protein. Lysosomal. May be localized in the endoplasmic reticulum.</p>

画像



ab60036 at 1/50 dilution staining ABCB9 in Human hepatocarcinoma by Immunohistochemistry, Formalin-fixed, Paraffin-embedded tissue. A peroxidase-conjugated secondary antibody was used, followed by AEC staining.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors