

Anti-Apolipoprotein C1/Apo-C1 antibody ab7617

製品の概要

製品名	Anti-Apolipoprotein C1/Apo-C1 antibody
製品の詳細	Goat polyclonal to Apolipoprotein C1/Apo-C1
由来種	Goat
特異性	Typically less than 1% cross reactivity against other types of apoLipoprotein was detected by ELISA. This antibody reacts with human apoLipoprotein C-I and has negligible cross-reactivity with Type A-I, A-II, B, C-II, C-III, E and J apoLipoproteins.
アプリケーション	適用あり: Dot blot, ELISA, WB
種交差性	交差種: Human
免疫原	Full length native protein (purified) corresponding to Apolipoprotein C1/Apo-C1.
特記事項	This antibody has been used to determine that atherosclerotic lesions in the human aorta contain considerable amounts of lipoproteins. These lipoproteins were observed to be complexed with components of the extracellular matrix (especially LDL and proteoglycans). The role of these matrix-lipoprotein complexes is not entirely clear, however, animal models of atherosclerosis have shown that increased cellular proliferation and increased production of extracellular matrix components occur following injury to the intimal layer of the aorta. This product was previously labelled as Apolipoprotein C1

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.01% Sodium azide Constituents: 4.7625% Sodium borate, 0.146% EDTA, 0.435% Sodium chloride
精製度	Immunogen affinity purified
特記事項 (精製)	This product has been prepared by immunoaffinity chromatography using immobilized antigens followed by extensive cross-adsorption against other apoLipoproteins and human serum proteins to remove any unwanted specificities.
ポリモノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab7617の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Dot blot		1/5000 - 1/10000.
ELISA		1/4000 - 1/8000.
WB		1/1000.

ターゲット情報

機能	Appears to modulate the interaction of APOE with beta-migrating VLDL and inhibit binding of beta-VLDL to the LDL receptor-related protein. Binds free fatty acids and reduces their intracellular esterification.
組織特異性	Synthesized mainly in liver and to a minor degree in intestine. Secreted in plasma.
配列類似性	Belongs to the apolipoprotein C1 family.
細胞内局在	Secreted.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors