

Anti-Aryl hydrocarbon Receptor antibody [RPT9] - CHIP Grade ab2769

★★★★☆ [9 Abreviews](#) [44 References](#) [画像数 11](#)

製品の概要

製品名	Anti-Aryl hydrocarbon Receptor antibody [RPT9] - CHIP Grade
製品の詳細	Mouse monoclonal [RPT9] to Aryl hydrocarbon Receptor - CHIP Grade
由来種	Mouse
特異性	This antibody specifically immunoprecipitates a single ~95 kDa protein representing AHR from Hepa 1 cytosol. Immunohistochemical staining of AHR in rat liver results in strong cytoplasmic and some nuclear staining.
アプリケーション	適用あり: ICC/IF, ELISA, WB, IHC-Fr, ChIP, IHC-P, IP, Flow Cyt
種交差性	交差種: Mouse, Rat, Human 交差が予測される動物種: Rabbit 
免疫原	Recombinant fragment corresponding to Mouse Aryl hydrocarbon Receptor aa 12-31. Sequence: R(12)KRRKP(17) V(22)KPIPAEGIK(31) Run BLAST with Run BLAST with
ポジティブ・コントロール	rat liver tissue sections Hepa 1 cytosolic lysate

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.05% Sodium azide
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	RPT9
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab2769の使用に適用されます


アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF	★★★★★ (1)	1/10 - 1/100.
ELISA		Use at an assay dependent concentration.
WB	★★★★★ (2)	1/500 - 1/2000. PubMed: 21266776 Albeit some customers have been successful using Ab2769 for western blots with endogenous samples, we recommend using ab190797 for this application.
IHC-Fr		Use at an assay dependent concentration. PubMed: 23036591
ChIP	★★★★★ (2)	1/100.
IHC-P	★★★★★ (2)	Use at an assay dependent concentration.
IP	★★★★★ (1)	Use at an assay dependent concentration.
Flow Cyt	★★★★★ (1)	Use 1-5µg for 10 ⁶ cells. ab18392 - Mouse monoclonal IgG3, is suitable for use as an isotype control with this antibody.

ターゲット情報


機能	Ligand-activated transcriptional activator. Binds to the XRE promoter region of genes it activates. Activates the expression of multiple phase I and II xenobiotic chemical metabolizing enzyme genes (such as the CYP1A1 gene). Mediates biochemical and toxic effects of halogenated aromatic hydrocarbons. Involved in cell-cycle regulation. Likely to play an important role in the development and maturation of many tissues.
組織特異性	Expressed in all tissues tested including blood, brain, heart, kidney, liver, lung, pancreas and skeletal muscle.
配列類似性	Contains 1 basic helix-loop-helix (bHLH) domain. Contains 1 PAC (PAS-associated C-terminal) domain. Contains 2 PAS (PER-ARNT-SIM) domains.
細胞内局在	Cytoplasm. Nucleus. Initially cytoplasmic; upon binding with ligand and interaction with a HSP90, it translocates to the nucleus.

画像




Western blot - Anti-Aryl hydrocarbon Receptor antibody [RPT9] - ChIP Grade (ab2769)

Western blot analysis of Aryl Hydrocarbon Receptor was performed by loading 40 ug of HEK293 lysate overexpressing Aryl Hydrocarbon Receptor (right lane) or empty vector control (left lane) and 10ul of a Prestained Protein Ladder onto a 4-20% Tris-HCl polyacrylamide gel. Proteins were transferred to a PVDF membrane and blocked with 5% BSA/TBST for at least 1 hour. The membrane was probed with an Aryl Hydrocarbon Receptor monoclonal antibody (ab2769) at a dilution of 1:1000 overnight at 4°C on a rocking platform then washed in TBS-0.1% Tween-20 and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:20,000 for 1 hour. Chemiluminescent detection was performed.




Immunocytochemistry/ Immunofluorescence - Anti-Aryl hydrocarbon Receptor antibody [RPT9] - ChIP Grade (ab2769)

Immunocytochemistry/Immunofluorescence analysis of Aryl Hydrocarbon Receptor shows staining in A2058 cells. Aryl Hydrocarbon Receptor (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were incubated without (control) or with ab2769 (1:20) overnight at 4°C, washed with PBS and incubated with a DyLight-488 conjugated secondary antibody. Images were taken at 60X magnification.




Immunocytochemistry/ Immunofluorescence - Anti-Aryl hydrocarbon Receptor antibody [RPT9] - ChIP Grade (ab2769)

Immunocytochemistry/Immunofluorescence analysis of Aryl Hydrocarbon Receptor shows staining in HeLa cells. Aryl Hydrocarbon Receptor (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were incubated without (control) or with ab2769 (1:20) overnight at 4°C, washed with PBS and incubated with a DyLight-488 conjugated secondary antibody. Images were taken at 60X magnification.




Immunocytochemistry/ Immunofluorescence - Anti-Aryl hydrocarbon Receptor antibody [RPT9] - ChIP Grade (ab2769)

Immunocytochemistry/Immunofluorescence analysis of Aryl Hydrocarbon Receptor shows staining in U251 cells. Aryl Hydrocarbon Receptor (green), F-Actin staining with Phalloidin (red) and nuclei with DAPI (blue) is shown. Cells were grown on chamber slides and fixed with formaldehyde prior to staining. Cells were incubated without (control) or with ab2769 (1:20) overnight at 4°C, washed with PBS and incubated with a DyLight-488 conjugated secondary antibody. Images were taken at 60X magnification.



Immunocytochemistry/ Immunofluorescence - Anti-Aryl hydrocarbon Receptor antibody [RPT9] - ChIP Grade (ab2769)

ICC/IF image of ab2769 stained HepG2 cells. The cells were methanol fixed (5 min) and then incubated in 1%BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab2769, in a 1/200 dilution) overnight at +4°C. The secondary antibody (green) was Alexa Fluor® 488 goat anti-mouse IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1h. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43µM.



ChIP - Anti-Aryl hydrocarbon Receptor antibody
[RPT9] - ChIP Grade (ab2769)

Image courtesy of an anonymous Abreview.

This was a trial experiment to evaluate the association of AhR to the ABCG2 promoter using ab2769 at a 1/100 dilution for the IP (ChIP assay). Semiquantitative PCR was performed to evaluate the relative association of AhR with the proximal ABCG2 promoter in a S1 colon cancer cell line without treatment, or treated with depsiptide (10 ng/mL 24h) or benzo(a)pyrene (10uM 24h).

Cross-linking (X-ChIP) - 10 Mins 0 Secs

Lane 1: DNA ladder (100 bp from promega)

Inputs (lanes 2-4): S1 no treatment - serial dilution

Inputs (lanes 5-7): S1 treated with depsiptide - serial dilution

Inputs (lanes 8-10): S1 treated with benzo(a)pyrene - serial dilution

AhR ChIP (lane 11-13): use 1:1 diluted immunoprecipitate for PCR


Lane 11=S1 no treatment lane 12=S1 depsiptide 10ng/mL lane

13= S1 benzo(a)pyrene 10uM

AhR ChIP (lane 14-16): use 1:2 diluted immunoprecipitate for PCR

same order as lanes 11-13


Lane 17: H2O control for PCR.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Aryl hydrocarbon Receptor antibody [RPT9] - ChIP Grade (ab2769)


This image is courtesy of an anonymous Abreview

ab2769 staining Aryl hydrocarbon Receptor in Mouse liver tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and antigen retrieval was by heat mediation in a citrate buffer. Samples were incubated with primary antibody (1/750) for 1 hour. Ab98784 (1/500) was used as the secondary antibody. Background staining due to secondary with positive stainind seen in the cytoplasm of the hepatocytes




Flow Cytometry - Anti-Aryl hydrocarbon Receptor antibody [RPT9] - ChIP Grade (ab2769)

Flow cytometry analysis of Aryl Hydrocarbon Receptor showing positive staining in the nucleus and cytoplasm of NIH/3T3 cells compared to an isotype control (blue). Cells were harvested and adjusted to a concentration of $1-5 \times 10^6$ cells/ml. Cells were then fixed with 2% paraformaldehyde and washed with PBS. Cells were penetrated by dropping the supernatant and adding 90% methanol followed by incubation for 10 minutes at room temperature. Cells were blocked with a 2% solution of BSA-PBS for 30 min at room temperature and incubated with ab2769 at 1:50 for 60 min at room temperature. Cells were then incubated for 40 min at room temperature in the dark using a Dylight 488-conjugated goat anti-mouse IgG (H+L) secondary antibody and re-suspended in PBS for FACS analysis.




Flow Cytometry - Anti-Aryl hydrocarbon Receptor antibody [RPT9] - ChIP Grade (ab2769)

Flow cytometry analysis of Aryl Hydrocarbon Receptor showing positive staining in the nucleus and cytoplasm of PC-3 cells compared to an isotype control (blue). Cells were harvested and adjusted to a concentration of $1-5 \times 10^6$ cells/ml. Cells were then fixed with 2% paraformaldehyde and washed with PBS. Cells were penetrated by dropping the supernatant and adding 90% methanol followed by incubation for 10 minutes at room temperature. Cells were blocked with a 2% solution of BSA-PBS for 30 min at room temperature and incubated with ab2769 at 1:50 for 60 min at room temperature. Cells were then incubated for 40 min at room temperature in the dark using a Dylight 488-conjugated goat anti-mouse IgG (H+L) secondary antibody and re-suspended in PBS for FACS analysis.



Flow cytometry analysis of Aryl Hydrocarbon Receptor showing positive staining in the nucleus and cytoplasm of HeLa cells compared to an isotype control (blue). Cells were harvested and adjusted to a concentration of $1-5 \times 10^6$ cells/ml. Cells were then fixed with 2% paraformaldehyde and washed with PBS. Cells were penetrated by dropping the supernatant and adding 90% methanol followed by incubation for 10 minutes at room temperature. Cells were blocked with a 2% solution of BSA-PBS for 30 min at room temperature and incubated with ab2769 at 1:50 for 60 min at room temperature. Cells were then incubated for 40 min at room temperature in the dark using a DyLight 488-conjugated goat anti-mouse IgG (H+L) secondary antibody and re-suspended in PBS for FACS analysis.



Overlay histogram showing HEK293 cells stained with ab2769 (red line). The cells were fixed with 100% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab2769, 1/20 dilution) for 30 min at 22°C. The secondary antibody used was a goat **anti mouse-DyLight® 488** (IgG H+L) (**ab96879**) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] (**ab91353**, 2µg/ 1×10^6 cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in HEK293 cells fixed with 4% paraformaldehyde (10 min)/permeabilized in 0.1% PBS-Tween used under the same conditions.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors