

Product datasheet

Anti-EGF antibody ab94898

製品の概要

製品名	Anti-EGF antibody
製品の詳細	Rabbit polyclonal to EGF
由来種	Rabbit
アプリケーション	適用あり: ELISA, WB, Indirect ELISA
種交差性	交差種: Human
免疫原	Recombinant Human EGF

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at 4°C (up to 6 months). Store at -20°C.
バッファー	Preservative: None Constituents: PBS, pH 7.2
精製度	Immunogen affinity purified
特記事項 (精製)	ab94898 was purified by affinity chromatography employing immobilized Human EGF matrix
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab94898** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
ELISA		1/0.5 - 1/2.
WB		Use a concentration of 0.1 - 0.2 µg/ml. Predicted molecular weight: 134 kDa. Detection limit is 1.5-3.0 ng, under either reducing or non-reducing conditions

アプリ ケー ション	Abreviews	特記事項
Indirect ELISA		Use a concentration of 0.5 - 2 µg/ml. To detect IL-10 by indirect ELISA (using 100 µl/well antibody solution) a concentration of 0.5 - 2.0 µg/ml of this antibody is required. This antigen affinity purified antibody, in conjunction with compatible secondary reagents, allows the detection of at least 0.2 - 0.4 ng/well of recombinant rIL-10.

ターゲット情報

機能	EGF stimulates the growth of various epidermal and epithelial tissues in vivo and in vitro and of some fibroblasts in cell culture. Magnesiotropic hormone that stimulates magnesium reabsorption in the renal distal convoluted tubule via engagement of EGFR and activation of the magnesium channel TRPM6. Can induce neurite outgrowth in motoneurons of the pond snail <i>Lymnaea stagnalis</i> in vitro (PubMed:10964941).
組織特異性	Expressed in kidney, salivary gland, cerebrum and prostate.
関連疾患	Hypomagnesemia 4
配列類似性	Contains 9 EGF-like domains. Contains 9 LDL-receptor class B repeats.
翻訳後修飾	O-glycosylated with core 1-like and core 2-like glycans. It is uncertain if Ser-954 or Thr-955 is O-glycosylated. The modification here shows glycan heterogeneity: HexHexNAc (major) and Hex2HexNAc2 (minor).
細胞内局在	Membrane.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors