


Product datasheet

Anti-GARNL3 antibody ab122015

画像数 3

製品の概要

| | |
|--------------|--|
| 製品名 | Anti-GARNL3 antibody |
| 製品の詳細 | Rabbit polyclonal to GARNL3 |
| 由来種 | Rabbit |
| アプリケーション | 適用あり: WB, ICC/IF, IHC-P |
| 種交差性 | 交差種: Human 交差が予測される動物種: Mouse  |
| 免疫原 | amino acids 737-828 (LVQPSASDFQ FCWNQAPYAI VCAFPYLLAF TTDSMEIRLV VNGNLVHTAV VPQLQLVASR SDIYFTATAA VNEVSSGGSS KGASARNSPQ TP) of Human GARNL3 (Q5VVW2). Run BLAST with ExPASy Run BLAST with NCBI |
| ポジティブ・コントロール | Human nasopharynx tissue. |

製品の特性

| | |
|--------|--|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle. |
| バッファー | pH: 7.20 Preservative: 0.02% Sodium azide Constituents: 59% PBS, 40% Glycerol |
| 精製度 | Immunogen affinity purified |
| ポリ/モノ | ポリクローナル |
| アイソタイプ | IgG |

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab122015** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|---|
| WB | | Use at an assay dependent concentration. |
| ICC/IF | | Use a concentration of 1 - 4 µg/ml. Recommend PFA Fixation and Triton X-100 treatment |
| IHC-P | | 1/20 - 1/50. Perform heat mediated antigen retrieval via the pressure cooker method before commencing with IHC staining protocol. |

ターゲット情報

配列類似性


Belongs to the GARNL3 family.
Contains 1 CNH domain.
Contains 1 Rap-GAP domain.

画像




Immunofluorescent staining of Human cell line U-251MG shows positivity in vesicles. Recommended concentration of ab122015 1-4 µg/ml. Cells treated with PFA/Triton X-100.

Immunocytochemistry/ Immunofluorescence - Anti-GARNL3 antibody (ab122015)



ab122015 staining GARNL3 in paraffin-embedded Human nasopharynx tissue by Immunohistochemistry.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-GARNL3 antibody (ab122015)



Western blot - Anti-GARNL3 antibody (ab122015)

Lane 1: Marker [kDa] 250; 130; 95; 72; 55; 36; 28; 17; 10
Lane 2: Negative control (vector only transfected HEK293T lysate)
Lane 3: Over-expression Lysate (Co-expressed with a C-terminal myc-DDK tag (~3.1 kDa) in mammalian HEK293T cells. Lane 1: Marker [kDa] 250; 130; 95; 72; 55; 36; 28; 17; 10
Lane 2: Negative control (vector only transfected HEK293T lysate)
Lane 3: Over-expression Lysate (Co-expressed with a C-terminal myc-DDK tag (~3.1 kDa) in mammalian HEK293T cells.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors