

Product datasheet

Anti-GNAT2 antibody ab97501

★★★★☆ 1 Abreviews 2 References 画像数 1

製品の概要

製品名	Anti-GNAT2 antibody
製品の詳細	Rabbit polyclonal to GNAT2
由来種	Rabbit
アプリケーション	適用あり: WB
種交差性	交差種: Mouse, Rat
免疫原	Synthetic peptide conjugated to KLH derived from within residues 150 - 250 of Mouse GNAT2.Immunogen の所有権に関して
ポジティブ・コントロール	This antibody gave a positive signal in both Rat and Mouse Retina tissue lysates.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.02% Sodium Azide Constituents: 1% BSA, PBS, pH 7.4
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab97501** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB	★★★★☆	Use a concentration of 1 µg/ml. Detects a band of approximately 54 kDa (predicted molecular weight: 40 kDa).

ターゲット情報

機能	Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems. Transducin is an amplifier and one of the transducers of a visual impulse that performs the coupling between rhodopsin and cGMP-phosphodiesterase.
組織特異性	Retinal rod outer segment.
関連疾患	Defects in GNAT2 are the cause of achromatopsia type 4 (ACHM4) [MIM:139340]. Achromatopsia is an autosomal recessively inherited visual disorder that is present from birth and that features the absence of color discrimination.
配列類似性	Belongs to the G-alpha family. G(i/o/t/z) subfamily.

画像



Western blot - Anti-GNAT2 antibody (ab97501)

All lanes : Anti-GNAT2 antibody (ab97501) at 1 µg/ml

Lane 1 : Mouse Retina Tissue Lysate

Lane 2 : Rat Retina Tissue Lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) preadsorbed (ab97080) at 1/5000 dilution

Developed using the ECL technique.

Performed under reducing conditions.

Predicted band size: 40 kDa

Observed band size: 54 kDa

Exposure time: 1 minute

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors