

Product datasheet

Anti-Glutathione Peroxidase 1 antibody ab22604

KO 評価済

★★★★★ 23 Abreviews 60 References 画像数 6

製品の概要

製品名	Anti-Glutathione Peroxidase 1 antibody
製品の詳細	Rabbit polyclonal to Glutathione Peroxidase 1
由来種	Rabbit
アプリケーション	適用あり: IHC-P, WB, ICC/IF, IHC-Fr
種交差性	交差種: Mouse, Rat, Cow, Human
免疫原	Synthetic peptide conjugated to KLH derived from within residues 150 to the C-terminus of Human Glutathione Peroxidase 1. Immunogen の所有権に関して (Peptide available as ab25301 .)
ポジティブ・コントロール	This antibody gave a positive signal in Human, Rat and mouse liver tissue lysates.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.02% Sodium Azide Constituents: 1% BSA, PBS, pH 7.4
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab22604** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.


アプリケーション	Abreviews	特記事項
IHC-P	★★★★★	1/300 - 1/1000. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

アプリケーション	Abreviews	特記事項
WB	★★★★☆	Use a concentration of 1 µg/ml. Detects a band of approximately 22 kDa (predicted molecular weight: 22 kDa).
ICC/IF	★★★★★	Use a concentration of 5 µg/ml.
IHC-Fr	★★★★★	Use at an assay dependent concentration.

ターゲット情報

機能	Protects the hemoglobin in erythrocytes from oxidative breakdown.
配列類似性	Belongs to the glutathione peroxidase family.
細胞内局在	Cytoplasm.


画像



Immunocytochemistry/ Immunofluorescence - Anti-Glutathione Peroxidase 1 antibody (ab22604)

This image is courtesy of an Abreview submitted by Carl Hobbs

ab22604 staining mouse Cor-1 cells by ICC/IF (Immunocytochemistry/immunofluorescence). Cells were fixed with formaldehyde, permeabilized with TBS/BSA/azide/0.1%Tween 20 and blocked with 1% BSA for 10 minutes at RT. Samples were incubated with primary antibody (1/300) for 2 hours. An diluted (1/1000) Alexa Fluor® 594-conjugated Goat anti-rabbit IgG polyclonal was used as the secondary antibody.



Western blot - Anti-Glutathione Peroxidase 1 antibody (ab22604)

Lane 1: Wild-type HAP1 cell lysate (20 µg)

Lane 2: GPX1 knockout HAP1 cell lysate (20 µg)


Lane 3: THP1 cell lysate (20 µg)

Lane 4: HL60 cell lysate (20 µg)

Lanes 1 - 4: Merged signal (red and green).

Green - ab22604 observed at 24 kDa. Red - loading control, [ab8245](#), observed at 37 kDa.

ab22604 was shown to recognize Glutathione Peroxidase 1 in wild-type HAP1 cells along with additional cross-reactive bands. No band was observed when Glutathione Peroxidase 1 knockout samples were examined. Wild-type and Glutathione Peroxidase 1 knockout samples were subjected to SDS-PAGE. ab22604 at a concentration of 1 µg/ml and [ab8245](#) (loading control to GAPDH) diluted at 1/2000 were incubated overnight at 4°C. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1/10,000 dilution for 1 hour at room temperature before imaging.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Glutathione Peroxidase 1 antibody (ab22604)

Image courtesy of [Human Protein Atlas](#)


ab22604 staining Glutathione Peroxidase 1 in Human kidney. The paraffin embedded human kidney tissue was incubated with ab22604 (1/375 dilution) for 30 mins at room temperature. Antigen retrieval was performed by heat induction in citrate buffer pH 6. Ab22604 was tested in a tissue microarray (TMA) containing a wide range of normal and cancer tissues as well as a cell microarray consisting of a range of commonly used, well characterised human cell lines.

Further results for this antibody can be found at www.proteinatlas.org



Immunocytochemistry/ Immunofluorescence - Anti-Glutathione Peroxidase 1 antibody (ab22604)

ICC/IF image of ab22604 stained human HepG2 cells. The cells were methanol fixed (5 min), permabilised in 0.1% PBS-Tween (20 min) and incubated with the antibody (ab22604, 5μg/ml) for 1h at room temperature. 1%BSA / 10% normal goat serum / 0.3M glycine was used to block non-specific protein-protein interactions. The secondary antibody (green) was Alexa Fluor® 488 goat anti-rabbit IgG (H+L) used at a 1/1000 dilution for 1h. Alexa Fluor® 594 WGA was used to label plasma membranes (red). DAPI was used to stain the cell nuclei (blue).



Western blot - Anti-Glutathione Peroxidase 1 antibody (ab22604)

All lanes : Anti-Glutathione Peroxidase 1 antibody (ab22604) at 1 µg/ml (Blocked with 3% Milk)

Lane 1 : Liver (Human) Tissue Lysate

Lane 2 : Liver (Mouse) Tissue Lysate

Lane 3 : Liver (Rat) Tissue Lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) preadsorbed at 1/50000 dilution

Developed using the ECL technique.

Performed under reducing conditions.


Predicted band size: 22 kDa

Observed band size: 22 kDa

Additional bands at: 190 kDa, 55 kDa, 65 kDa. We are unsure as to the identity of these extra bands.

Exposure time: 4 minutes

This blot was produced using a 4-12% Bis-tris gel under the MES buffer system. The gel was run at 200V for 35 minutes before being transferred onto a Nitrocellulose membrane at 30V for 70 minutes. The membrane was then blocked for an hour using 3% milk before being incubated with ab22604 overnight at 4°C. Antibody binding was detected using an anti-rabbit antibody conjugated to HRP, and visualised using ECL development solution [ab133406](#).



Western blot - Anti-Glutathione Peroxidase 1 antibody (ab22604)

All lanes : Anti-Glutathione Peroxidase 1 antibody (ab22604) at 1 µg/ml

Lane 1 : Human liver normal tissue lysate

Lane 2 : Liver (Mouse) Tissue Lysate (ab7935)

Lane 3 : Human liver normal tissue lysate with Human Glutathione Peroxidase 1 peptide (ab25301) at 1 µg/ml

Lane 4 : Liver (Mouse) Tissue Lysate (ab7935) with Human Glutathione Peroxidase 1 peptide (ab25301) at 1 µg/ml

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat polyclonal to Rabbit IgG at 1/10000 dilution

Performed under reducing conditions.

Predicted band size: 22 kDa

Observed band size: 22,24 kDa

ab22604 detects a band of approximately 22 kDa in human liver lysate. The band detected in the mouse liver lysate runs slightly higher (~24kDa). We believe that these bands correspond to Glutathione Peroxidase 1 as they are both blocked by the addition of the immunizing peptide (ab25301).

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish

- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors