

## Product datasheet

# Goat Anti-Rabbit IgG H&L (HRP) preadsorbed ab7090

★★★★★ 2 Abreviews 28 References 画像数 2

### 製品の概要

製品名	Goat Anti-Rabbit IgG H&L (HRP) preadsorbed
由来種	Goat
ターゲット生物種	Rabbit
アプリケーション	<b>適用あり:</b> ICC/IF, IHC-P, WB, ELISA, Immunomicroscopy, Dot blot, IHC-Fr
吸着処理血清	Chicken, Cow, Goat, Guinea pig, Hamster, Horse, Human, Mouse, Rat, Sheep <a href="#">more details</a>
免疫原	Full length native Rabbit IgG (purified).
標識	HRP

### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C.
バッファー	pH: 7.20 Preservative: 0.01% Gentamicin sulphate Constituents: 0.42% Potassium phosphate, 0.87% Sodium chloride, 1% BSA
精製度	Affinity purified
特記事項 (精製)	This product was prepared from monospecific antiserum by immunoaffinity chromatography using Rabbit IgG coupled to agarose beads.
特記事項 (標識)	Horseradish Peroxidase (HRP)
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

### アプリケーション


Our [Abpromise guarantee](#) covers the use of **ab7090** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
ICC/IF		1/1000 - 1/5000.

アプリケーション	Abreviews	特記事項
IHC-P	★★★★☆	Use at an assay dependent concentration.
WB	★★★★★	Use at an assay dependent concentration.
ELISA		1/20000 - 1/80000.
Immunomicroscopy		Use at an assay dependent concentration.
Dot blot		Use at an assay dependent concentration.
IHC-Fr		Use at an assay dependent concentration.


## 画像



Immunohistochemical analysis of healthy and colitis mouse colon sections (untreated and treated with enoxaparin) labeling claudin-4 with [ab15104](#) at 1/200. [ab7090](#), anti-rabbit immunoglobulin G conjugated to horseradish peroxidase (HRP) at 1/300 was used as the secondary antibody. [ab64238](#) at 1/50 was used to develop the histological signal.

Antigen retrieval was performed by incubating the sections for 10 minutes at 97°C in 1 mM EDTA buffer, pH 8.0 or 10 mM citrate buffer, pH 6.0.

Control, C; untreated colitis, DSS; oral enoxaparin, OE; intraperitoneal injection of enoxaparin, IPE.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Goat Anti-Rabbit IgG H&L (HRP) preadsorbed (ab7090)

This image is courtesy of an anonymous Abreview.

**ab9986** staining osteoprotegerin in human skin tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Tissue was fixed with formaldehyde and blocked with 1% BSA and 5% normal goat serum in PBS for 30 minutes at 22°C; Samples were incubated with primary antibody (1/20) for 9 hours at 4°C. An HRP-conjugated goat anti-rabbit IgG H&L (ab7090) (1/300) was used as the secondary antibody.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors