

Product datasheet

Anti-ING3 antibody ab115470

★★★★★ 1 Abreviews [画像数 1](#)

製品の概要

製品名	Anti-ING3 antibody
製品の詳細	Goat polyclonal to ING3
特異性	ab115470 is specific for isoform 1 of ING3.
アプリケーション	<b>適用あり:</b> WB, ELISA, IHC-P, ICC/IF
種交差性	<b>交差種:</b> Human <b>交差が予測される動物種:</b> Mouse, Rat, Horse, Chicken, Cow, Dog, Pig 
免疫原	Synthetic peptide: ADSRSGRKSKN conjugated to KLH, corresponding to amino acids 294-304 of Human ING3. <a href="#">Run BLAST with</a> <a href="#">Run BLAST with</a>
ポジティブ・コントロール	Human brain (cortex) tissue.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 0.27% Potassium phosphate, 0.88% Sodium chloride
精製度	Protein G purified
ポリモノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab115470** in the following tested applications.  
The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB	★★★★☆	1/200 - 1/2000. Predicted molecular weight: 47 kDa.
ELISA		1/10000 - 1/40000.
IHC-P		Use a concentration of 2.5 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
ICC/IF		Use at an assay dependent dilution.

## ターゲット情報

### 関連性


**Function:** Component of the NuA4 histone acetyltransferase (HAT) complex which is involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A. This modification may both alter nucleosome - DNA interactions and promote interaction of the modified histones with other proteins which positively regulate transcription. This complex may be required for the activation of transcriptional programs associated with oncogene and proto-oncogene mediated growth induction, tumor suppressor mediated growth arrest and replicative senescence, apoptosis, and DNA repair. NuA4 may also play a direct role in DNA repair when directly recruited to sites of DNA damage. Component of a SWR1-like complex that specifically mediates the removal of histone H2A.Z/H2AFZ from the nucleosome.

**Tissue specificity:** Expressed in brain, heart, kidney, liver, lung, ovaries, placenta, prostate, skeletal muscle, small intestine, spleen, testis and thymus. **Disease:** Squamous cell carcinoma of the head and neck **Similarity:** Belongs to the ING family. Contains 1 PHD-type zinc finger. **Domain:** The PHD-type zinc finger mediates the binding to H3K4me3.

### 細胞内局在

Nuclear

## 画像



ab115470 at 2.5ug/ml staining ING3 in Formalin-fixed, Paraffin-embedded Human brain (cortex) tissue by Immunohistochemistry, followed by biotinylated secondary antibody, alkaline phosphatase-streptavidin and chromogen.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ING3 antibody (ab115470)

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <http://www.abcam.co.jp/abpromise> or contact our technical team.

## **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors