

Product datasheet

Anti-Nova1 antibody ab77594

★★★★☆ 1 Abreviews 3 References 画像数 3

製品の概要

製品名	Anti-Nova1 antibody
製品の詳細	Goat polyclonal to Nova1
由来種	Goat
特異性	ab77594 is expected to recognise all three reported isoforms (NP_002506.2 ; NP_006480.2 ; NP_006482.1).
アプリケーション	適用あり: WB, IHC-P
種交差性	交差種: Human 交差が予測される動物種: Mouse, Rat, Cynomolgus monkey
免疫原	Synthetic peptide: REMPQNVAKTEPVS , corresponding to internal sequence amino acids 124-137 of Human Nova1 (NP_002506.2 ; NP_006480.2 ; NP_006482.1). Run BLAST with Run BLAST with
ポジティブ・コントロール	Human breast cancer lysate. Transfected HEK293 transiently expressing Nova1. Human cerebral cortex tissue sections.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	Preservative: 0.02% Sodium Azide Constituents: 0.5% BSA, Tris buffered saline, pH 7.3
精製度	Immunogen affinity purified
特記事項(精製)	ab77594 was purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab77594** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
WB	★★★★☆	Use a concentration of 0.01 - 0.03 µg/ml. Detects a band of approximately 52 kDa (predicted molecular weight: 52 kDa).
IHC-P		Use a concentration of 2 - 4 µg/ml.

ターゲット情報

機能	May regulate RNA splicing or metabolism in a specific subset of developing neurons.
組織特異性	Brain.
配列類似性	Contains 3 KH domains.
細胞内局在	Nucleus.

画像




Anti-Nova1 antibody (ab77594) at 0.01 µg/ml
+ Human Breast Cancer lysate (in RIPA buffer)
at 35 µg


Developed using the ECL technique.

Predicted band size: 52 kDa

Observed band size: 52 kDa



ab77594 at 2.5µg/ml staining Nova1 in human cerebral cortex tissue by Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections). Steamed antigen retrieval was performed with citrate buffer pH 6, AP-staining.



All lanes : Anti-Nova1 antibody (ab77594) at 0.03 µg/ml

Lane 1 : Lysate prepared from MOCK transfection.

Lane 2 : Lysate prepared from transfected HEK293 cells, transiently expressing Nova1

Predicted band size: 52 kDa

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors