

Product datasheet

Anti-Nucleoporin p62 antibody ab96134

★★★★★ 4 Abreviews 6 References 画像数 6

製品の概要

| | |
|--------------|--|
| 製品名 | Anti-Nucleoporin p62 antibody |
| 製品の詳細 | Rabbit polyclonal to Nucleoporin p62 |
| 由来種 | Rabbit |
| アプリケーション | 適用あり: IHC-P, WB, ICC/IF |
| 種交差性 | 交差種: Mouse, Rat, Human 交差が予測される動物種: Cow, Zebrafish, Rhesus monkey ▲ |
| 免疫原 | Recombinant fragment corresponding to a region within amino acids 294 - 502 of Human Nucleoporin p62 (NP_057637). |
| ポジティブ・コントロール | WB: A431, H1299, PC-12, Rat2, Neuro-2a, GL261, C8D30, NIH/3T3, BCL-1, RAW 264.7 and C2C12 whole cell lysate. ICC/IF: A431 cells. IHC-P: Mouse fore brain tissue. |

法規制情報

医薬用外毒物

製品の特性

| | |
|--------|---|
| 製品の状態 | Liquid |
| 保存方法 | Shipped at 4°C. Store at +4°C short term (1-2 weeks). Store at -20°C or -80°C. Avoid freeze / thaw cycle. |
| バッファー | Preservative: 0.01% Thimerosal (merthiolate) Constituents: 10% Glycerol, 0.1M Tris, 0.1M Glycine, pH 7.0 |
| 精製度 | Immunogen affinity purified |
| ポリ/モノ | ポリクローナル |
| アイソタイプ | IgG |

アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab96134** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|---|
| IHC-P | | 1/100 - 1/1000. |
| WB | ★★★★★ | 1/500 - 1/3000. Predicted molecular weight: 53 kDa. |
| ICC/IF | ★★★★★ | 1/100 - 1/200. |

ターゲット情報


関連性

The nuclear pore complex is a structure that extends across the nuclear envelope and regulates the flow of macromolecules between the cytoplasm and the nucleus. Nucleoporins are the main components of the nuclear pore complex in eukaryotic cells. Nup 62 is localized to the nuclear pore central plug. This protein associates with the importin alpha/beta complex which is involved in the import of proteins containing nuclear localization signals. There are multiple transcript variants of this gene, however, they encode a single protein isoform. This protein undergoes post-translational modifications. It contains about 10 N-acetylglucosamine side chain sites.

細胞内局在

Nucleus; nuclear pore complex. Cytoplasm; cytoskeleton; spindle pole. Note: Central region of the nuclear pore, within the transporter. During mitotic cell division, it associates with the poles of the mitotic spindle.

画像



Western blot - Anti-Nucleoporin p62 antibody (ab96134)

All lanes : Anti-Nucleoporin p62 antibody (ab96134) at 1/1000 dilution

Lane 1 : Neuro-2a (Mouse neuroblastoma cell line) whole cell lysate

Lane 2 : GL261 whole cell lysate

Lane 3 : C8D30 whole cell lysate

Lane 4 : NIH/3T3 (mouse embryo fibroblast cell line) whole cell lysate

Lane 5 : BCL-1 (mouse B lymphocyte lymphoma cell line) whole cell lysate

Lane 6 : RAW 264.7 (Mouse macrophage cell line transformed with Abelson murine leukemia virus) whole cell lysate

Lane 7 : C2C12 (Mouse myoblast cell line) whole cell lysate


Lysates/proteins at 30 µg per lane.

Secondary

All lanes : HRP-conjugated anti-rabbit IgG


Predicted band size: 53 kDa

10% SDS-PAGE.



Immunocytochemistry/ Immunofluorescence - Anti-Nucleoporin p62 antibody (ab96134)

ab96134, at a 1/200 dilution, staining Nucleoporin p62 in paraformaldehyde fixed A431 by Immunofluorescence analysis. Lower image: merged with DNA probe.



Western blot - Anti-Nucleoporin p62 antibody
(ab96134)

All lanes : Anti-Nucleoporin p62 antibody
(ab96134) at 1/1000 dilution

Lane 1 : PC-12 (Rat adrenal gland
pheochromocytoma cell line) whole cell lysate

Lane 2 : Rat2 (Rat fibroblast cell line) whole
cell lysate


Lysates/proteins at 30 µg per lane.

Secondary

All lanes : HRP-conjugated anti-rabbit IgG


Predicted band size: 53 kDa

10% SDS-PAGE.



Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-Nucleoporin p62 antibody
(ab96134)

Paraffin-embedded mouse fore brain tissue
stained for Nucleoporin p62 using ab96134 at
1/500 dilution in immunohistochemical
analysis.



Western blot - Anti-Nucleoporin p62 antibody (ab96134)

All lanes : Anti-Nucleoporin p62 antibody (ab96134) at 1/1000 dilution


Lane 1 : A431 whole cell lysate

Lane 2 : H1299 whole cell lysate

Lysates/proteins at 30 µg per lane.

Predicted band size: 53 kDa

10% SDS PAGE



Immunocytochemistry/ Immunofluorescence - Anti-Nucleoporin p62 antibody (ab96134)

Immunofluorescence analysis of A431 cells, staining Nucleoporin p62 with ab96134 at 1/200 dilution, demonstrating punctate nuclear staining.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors