

## Product datasheet

# Recombinant Human Argininosuccinate Lyase protein ab113605

画像数 1

### 製品の概要

製品名	Recombinant Human Argininosuccinate Lyase protein
タンパク質長	Full length protein

### 製品の詳細

由来	Recombinant
由来	Escherichia coli

### アミノ酸配列

アクセッション番号 [P04424](#)

生物種 Human

配列 **MGSSHHHHHSSGLVPRGSHMAESGKLWGRFVGAVDPIMEKFNASIA Y  
DRHLWEVDVQGSKAYSRLGLEKAGLLTKAEMDQILHGLDKVAEEWAQGTK  
LNSNDEDIHTANERRLKEIGATAGKLHTGRSRNDQVVDLRLWMRQTCS  
TSLGLLWELIRTMVDRAEAERDVLFPGYTHLQRAQPIRWSHWILSHAVAL  
TRDSERLLEVRKRINVLPLGSGAIAGNPLGVDRELLRAELNFGAITLNSM  
DATSERDFVAEFLFASLCMTHLSRMAEDLILYCTKEFSFVQLSDAYSTG  
SSLMPQKKNPDSLELIRSKAGRVFGRCAGLLMTLKGLPSTYKDLQEDKE  
AVFEVSDTMSAVLQVATGVISTLQIHQENMGQALSPDMLATDLAYYLVRK  
GMPFRQAHEASGKAVFMAETKGVALNQLSLQELQTI SPLFSGDVICVWDY  
GHSVEQYGALGGTARSSVDWQIRQVRALLQAQQA**

分子量 54 kDa including tags

領域 1 to 464

タグ His tag N-Terminus

### 特性

Our [Abpromise guarantee](#) covers the use of **ab113605** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	SDS-PAGE
精製度	> 95 % SDS-PAGE. ab113605 was purified using conventional chromatography techniques.
製品の状態	Liquid

## 前処理および保存

### 保存方法および安定性

Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.

pH: 8.00

Constituents: 0.32% Tris HCl, 0.04% DTT, 10% Glycerol, 0.58% Sodium chloride

## 関連情報

### パスウェイ

Amino-acid biosynthesis; L-arginine biosynthesis; L-arginine from L-ornithine and carbamoyl phosphate: step 3/3.

Nitrogen metabolism; urea cycle; L-arginine and fumarate from (N(omega)-L-arginino)succinate: step 1/1.

### 関連疾患

Defects in ASL are the cause of arginosuccinicaciduria (ARGINSA) [MIM:207900].

Arginosuccinicaciduria is an autosomal recessive disorder of the urea cycle. The disease is characterized by mental and physical retardation, liver enlargement, skin lesions, dry and brittle hair showing trichorrhexis nodosa microscopically and fluorescing red, convulsions, and episodic unconsciousness.


### 配列類似性

Belongs to the lyase 1 family. Argininosuccinate lyase subfamily.

### 翻訳後修飾

Acetylation modifies enzyme activity in response to alterations of extracellular nutrient availability. Acetylation increased with trichostin A (TSA) or with nicotinamide (NAM). Glucose increases acetylation by about a factor of 3 with decreasing enzyme activity. Acetylation on Lys-288 is decreased on the addition of extra amino acids resulting in activation of enzyme activity.

## 画像



15% SDS-PAGE showing ab113605 (3 µg) at approximately 53.8 kDa.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors