

## Product datasheet

# Recombinant human VEGFC protein ab51968

### 1 References

#### 製品の概要

---

製品名	Recombinant human VEGFC protein
タンパク質長	Full length protein

#### 製品の詳細

---

由来	Recombinant
由来	Escherichia coli
アミノ酸配列	
生物種	Human

#### 特性

---

Our [Abpromise guarantee](#) covers the use of **ab51968** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

生理活性	Activity of VEGF-C is tested by the stimulation of VEGF-R3 phosphorylation in cells transfected with VEGF-R3. The expected ED <sub>50</sub> is > 200 ng/ml.
------	---

アプリケーション	SDS-PAGE
----------	----------

精製度	> 90 % SDS-PAGE. Endotoxin level is <0.1 ng per g of VEGF-C
-----	--

製品の状態	Lyophilised
-------	-------------

備考	Recombinant VEGF-C (Vascular Endothelial Growth Factor-C) is a 125 amino acid protein that lacks the N-terminal signal peptide
----	--

#### 前処理および保存

---

保存方法および安定性	Shipped at 4°C. Upon delivery aliquot. Store at -80°C. Avoid freeze / thaw cycle. This product is an active protein and may elicit a biological response in vivo, handle with caution.
再構成	reconstitution in 0.1% acetic acid to a concentration of 0.1-1.0 mg/ml.

## 関連情報

---

<b>機能</b>	Growth factor active in angiogenesis, and endothelial cell growth, stimulating their proliferation and migration and also has effects on the permeability of blood vessels. May function in angiogenesis of the venous and lymphatic vascular systems during embryogenesis, and also in the maintenance of differentiated lymphatic endothelium in adults. Binds and activates VEGFR-2 (KDR/FLK1) and VEGFR-3 (FLT4) receptors.
<b>組織特異性</b>	Spleen, lymph node, thymus, appendix, bone marrow, heart, placenta, ovary, skeletal muscle, prostate, testis, colon and small intestine and fetal liver, lung and kidney, but not in peripheral blood lymphocyte.
<b>配列類似性</b>	Belongs to the PDGF/VEGF growth factor family.
<b>翻訳後修飾</b>	Undergoes a complex proteolytic maturation which generates a variety of processed secreted forms with increased activity toward VEGFR-3, but only the fully processed form could activate VEGFR-2. VEGF-C first form an antiparallel homodimer linked by disulfide bonds. Before secretion, a cleavage occurs between Arg-227 and Ser-228 producing a heterotetramer. The next extracellular step of the processing removes the N-terminal propeptide. Finally the mature VEGF-C is composed mostly of two VEGF homology domains (VHDs) bound by non-covalent interactions.
<b>細胞内局在</b>	Secreted.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

## Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors