

## Product datasheet

### Anti-ZIC5 antibody ab115566

画像数 3

#### 製品の概要

製品名	Anti-ZIC5 antibody
製品の詳細	Goat polyclonal to ZIC5
由来種	Goat
アプリケーション	<b>適用あり:</b> ELISA, IHC-P
種交差性	<b>交差種:</b> Human
免疫原	Synthetic peptide: EIYGNPEVVRTIH with a Cysteine residue linker, corresponding to C terminal amino acids 651-663 of Human ZIC5 (NP_149123.2). <a href="#">Run BLAST with</a> <a href="#">Run BLAST with</a>
ポジティブ・コントロール	Human Prostate, Human Skin and Human Small Intestine tissues.

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: 99% Tris buffered saline, 0.5% BSA
精製度	Protein G purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

#### アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab115566** in the following tested applications.


The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
ELISA		1/128000.
IHC-P		Use a concentration of 3.75 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

## ターゲット情報


機能	Essential for neural crest development, converting cells from an epidermal fate to a neural crest cell fate. Binds to DNA.
配列類似性	Belongs to the GLI C2H2-type zinc-finger protein family. Contains 4 C2H2-type zinc fingers.
細胞内局在	Nucleus.

## 画像




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ZIC5 antibody (ab115566)

ab115566, at 3.75µg/ml, staining ZIC5 in Formalin-fixed, Paraffin-embedded Human Prostate tissue by Immunohistochemistry followed by biotinylated secondary antibody, alkaline phosphatase-streptavidin and chromogen.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ZIC5 antibody (ab115566)

ab115566, at 3.75µg/ml, staining ZIC5 in Formalin-fixed, Paraffin-embedded Human Skin tissue by Immunohistochemistry followed by biotinylated secondary antibody, alkaline phosphatase-streptavidin and chromogen.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ZIC5 antibody (ab115566)

ab115566, at 3.75µg/ml, staining ZIC5 in Formalin-fixed, Paraffin-embedded Human Small intestine tissue by Immunohistochemistry followed by biotinylated secondary antibody, alkaline phosphatase-streptavidin and chromogen. .

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors