

Product datasheet

Human lung tissue lysate (squamous cell carcinoma) - soluble fraction (male, 74 years) ab42216

製品の概要

製品名	Human lung tissue lysate (squamous cell carcinoma) - soluble fraction (male, 74 years)
特記事項	<p>Tissue: Lung Location: Left upper lobe. Diagnosis: Squamous cell carcinoma Stage: I T₂N₀M₀ Grade: II Gross findings: Tumor size 7 cm diameter, well demarcated. Cut section soft, hemorrhagic and necrotic. Ipsilateral hilar lymph nodes show hyperplasia. Microscopic findings: Tumor shows proliferation of malignant epithelial cell clusters. Epithelial cells form clusters with irregular, basophilic, large nuclei and predominant nucleoli. Cytoplasm is abundant and clear. Nuclear chromatin is coarse and irregularly distributed. Mitotic figures are evident. The surrounding stroma is hemorrhagic and infiltrated by large numbers of lymphocytes, plasma cells, eosinophils and neutrophils. Tumor necrosis and blood vessel invasion are revealed.</p> <p>This product belongs to a range of individual sets of lysates that were extracted from very specific tissues and are gender-, cancer subtype-, tumor grade- and cellular morphology-specific. They allow for consistency, reproducibility, and comparative studies from experiment to experiment, due to the existence of specific (matched) lysates that can be used as controls. Two extractions from each tissue were performed, extraction 1 yielding the soluble protein fraction and extraction 2 the insoluble protein fraction. Please find below a range of products that relate to ab42216: Extraction 1: soluble protein fraction ab42216 Human Lung Tumor Tissue lysate ab42215 Human Lung Normal Tissue lysate (matched) Extraction 2: insoluble protein fraction ab42218 Human Lung Tumor Tissue lysate ab42217 Human Lung Normal Tissue lysate (matched)</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped on Dry Ice. Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	<p>pH: 7.40</p> <p>Constituents: PBS, 0.0001% Pepstatin A, 0.018% Sodium orthovanadate, 0.004% Sodium fluoride, 0.0001% Aprotinin, 0.0001% Leupeptin, 0.25% Sodium deoxycholate, 0.017% PMSF, 0.1% SDS, 0.029% EDTA</p>
特記事項 (精製)	Tissue specimens are homogenized in modified RIPA buffer to obtain the soluble proteins, and centrifuged to clarify. The lysate solution may appear turbid at cold temperatures due to insolubility of buffer components. The solution should clear upon warming to room temperature.

Please note: All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors