

## Product datasheet

### Anti-ATG9A antibody [14F2 8B1] ab71795

#### 4 References

#### 製品の概要

製品名	Anti-ATG9A antibody [14F2 8B1]
製品の詳細	Armenian Hamster monoclonal [14F2 8B1] to ATG9A
由来種	Armenian hamster
アプリケーション	<b>適用あり:</b> ICC/IF, IHC-Fr, ELISA, WB, IP
種交差性	<b>交差種:</b> Mouse, Rat, Human
免疫原	Synthetic peptide: HPEPVPEEGSEDELPPQVHK (Human) <a href="#">Run BLAST with</a> <a href="#">Run BLAST with</a>
エピトープ	HPEPVPEEGSEDELPPQVHK
ポジティブ・コントロール	Endogenous protein, tagged over expressed protein and sRNA-depleted HEK 293 cells.
特記事項	The host species is Armenian hamster.

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	Preservative: None Constituents: PBS
精製度	Immunogen affinity purified
ポリ/モノ	モノクローナル
クローン名	14F2 8B1
ミエローマ	P3x63-Ag8.653
アイソタイプ	IgG

#### アプリケーション

Our [Abpromise guarantee](#) covers the use of **ab71795** in the following tested applications.

The application notes include recommended starting dilutions; optimal dilutions/concentrations should be determined by the end user.

アプリケーション	Abreviews	特記事項
ICC/IF		
IHC-Fr		
ELISA		
WB		
IP		

#### 追加情報

ELISA: Use at an assay dependent dilution.  
ICC/IF: Use at an assay dependent dilution.  
IHC-Fr: Use at an assay dependent dilution. Acetone fixed.  
IP: Use at an assay dependent dilution.  
WB: Use at an assay dependent dilution. Predicted molecular weight: 94 kDa.

Not yet tested in other applications.  
Optimal dilutions/concentrations should be determined by the end user.

#### ターゲット情報

##### 機能

Involved in autophagy and cytoplasm to vacuole transport (Cvt) vesicle formation. Plays a key role in the organization of the preautophagosomal structure/phagophore assembly site (PAS), the nucleating site for formation of the sequestering vesicle. Cycles between a juxta-nuclear trans-Golgi network compartment and late endosomes. Nutrient starvation induces accumulation on autophagosomes. Starvation-dependent trafficking requires ULK1, ATG13 and SUPT20H.

##### 配列類似性

Belongs to the ATG9 family.

##### 細胞内局在

Cytoplasmic vesicle, autophagosome membrane. Golgi apparatus, trans-Golgi network membrane. Late endosome membrane. Endoplasmic reticulum membrane. Under amino acid starvation or rapamycin treatment, redistributes from a juxtannuclear clustered pool to a dispersed peripheral cytosolic pool. The starvation-induced redistribution depends on ULK1, ATG13, as well as SH3GLB1.

**Please note:** All products are "FOR RESEARCH USE ONLY AND ARE NOT INTENDED FOR DIAGNOSTIC OR THERAPEUTIC USE"

#### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

## Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors