

Product datasheet

Anti-CD105 antibody [MEM-226] ab2529

KO 評価済

17 References 画像数 5

製品の概要

製品名	Anti-CD105 antibody [MEM-226]
製品の詳細	Mouse monoclonal [MEM-226] to CD105
由来種	Mouse
アプリケーション	適用あり: ICC/IF, Sandwich ELISA, Flow Cyt, WB
種交差性	交差種: Human
免疫原	Recombinant full length protein (Human). Expressed in vaccinia virus containing CD105 cDNA.
ポジティブ・コントロール	ICC/IF: HeLa cells. WB: Human colon tissue lysate. Flow Cyt: U937 cells.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.097% Sodium azide Constituent: PBS
精製度	Protein A purified
特記事項(精製)	Purified from TCS. Purity >95% by SDS-PAGE.
ポリ/モノ	モノクローナル
クローン名	MEM-226
アイソタイプ	IgG1

アプリケーション


The Abpromise guarantee [Abpromise保証は](#)、次のテスト済みアプリケーションにおけるab2529の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		Use a concentration of 10 µg/ml.
Sandwich ELISA		Use a concentration of 5 µg/ml. Can be paired for Sandwich ELISA with Rabbit polyclonal to CD105 (ab21224) . For sandwich ELISA, use this antibody as Capture at 5 µg/ml with Rabbit polyclonal to CD105 (ab21224) as Detection.
Flow Cyt		Use a concentration of 1 - 2 µg/ml. ab170190 - Mouse monoclonal IgG1, is suitable for use as an isotype control with this antibody.
WB		Use at an assay dependent concentration. Use under non reducing condition.

ターゲット情報

機能	Major glycoprotein of vascular endothelium. May play a critical role in the binding of endothelial cells to integrins and/or other RGD receptors.
組織特異性	Endoglin is restricted to endothelial cells in all tissues except bone marrow.
関連疾患	Defects in ENG are the cause of hereditary hemorrhagic telangiectasia type 1 (HHT1) [MIM:187300, 108010]; also known as Osler-Rendu-Weber syndrome 1 (ORW1). HHT1 is an autosomal dominant multisystemic vascular dysplasia, characterized by recurrent epistaxis, muco-cutaneous telangiectases, gastro-intestinal hemorrhage, and pulmonary (PAVM), cerebral (CAVM) and hepatic arteriovenous malformations; all secondary manifestations of the underlying vascular dysplasia. Although the first symptom of HHT1 in children is generally nose bleed, there is an important clinical heterogeneity.
細胞内局在	Membrane.

画像



Western blot - Anti-CD105 antibody [MEM-226] (ab2529)

All lanes : Anti-CD105 antibody [MEM-226] (ab2529) at 1/1000 dilution

Lane 1 : Wild-type HeLa whole cell lysate


Lane 2 : CD105 knockout HeLa whole cell lysate

Lane 3 : HUVEC whole cell lysate

Lysates/proteins at 20 µg per lane.

Lanes 1 - 3: Merged signal (red and green). Green - ab2529 observed at 70 kDa. Red - loading control, [ab181602](#), observed at 37 kDa.


ab2529 was shown to recognize ENG (Endoglin) in wild-type HeLa cells as signal was lost at the expected MW in ENG knockout cells. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and ENG knockout samples were subjected to SDS-PAGE. The membrane was blocked with 3% milk. Ab2529 and [ab181602](#) (Rabbit anti-GAPDH loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Mouse IgG H&L (IRDye® 800CW) preabsorbed [ab216772](#) and Goat anti-Rabbit IgG H&L (IRDye® 680RD) preabsorbed [ab216777](#) secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Flow Cytometry - Anti-CD105 antibody [MEM-226] (ab2529)

Overlay histogram showing U937 (Human histiocytic lymphoma cell line) cells stained with ab2529 (red line).


The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab2529, 1µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was DyLight[®] 488 goat anti-mouse IgG (H+L) (ab96879) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG1 [ICIGG1] (ab91353, 2µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in U937 cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Immunocytochemistry/ Immunofluorescence - Anti-CD105 antibody [MEM-226] (ab2529)


ICC/IF image of ab2529 stained HeLa (Human epithelial cell line from cervix adenocarcinoma) cells.

The cells were 4% formaldehyde fixed (10 min) and then incubated in 1% BSA / 10% normal goat serum / 0.3M glycine in 0.1% PBS-Tween for 1 hour to permeabilize the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody (ab2529, 10 µg/ml) overnight at +4°C. The secondary antibody (green) was ab69879, DyLight[®] 488 goat anti-mouse IgG (H+L) used at a 1/250 dilution for 1 hour. Alexa Fluor[®] 594 WGA was used to label plasma membranes (red) at a 1/200 dilution for 1 hour. DAPI was used to stain the cell nuclei (blue) at a concentration of 1.43 µM.



Sandwich ELISA - Anti-CD105 antibody [MEM-226] (ab2529)

Standard Curve for CD105 (Analyte: CD105 protein (ab54338)); dilution range 1pg/ml to 1ug/ml using Capture Antibody Mouse monoclonal [MEM-226] to CD105 (ab2529) at 5ug/ml and Detector Antibody Rabbit polyclonal to CD105 (ab21224) at 0.5ug/ml.



Western blot - Anti-CD105 antibody [MEM-226]
(ab2529)

Anti-CD105 antibody [MEM-226] (ab2529) at 5 µg/ml + Human colon tissue lysate at 10 µg

Secondary

Goat polyclonal to Mouse IgG - H&L - Pre-Adsorbed (HRP) at 1/3000 dilution

Observed band size: 80 kDa

Additional bands at: 45 kDa, 55 kDa. We are unsure as to the identity of these extra bands.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors