

# Anti-hnRNP A1 antibody ab4791

★★★★★ [1 Abreviews](#) [8 References](#) [画像数 5](#)

### 製品の概要

---

製品名	Anti-hnRNP A1 antibody
製品の詳細	Rabbit polyclonal to hnRNP A1
由来種	Rabbit
特異性	This antibody recognises hnRNPA1 (39kDa) in Western blots.
アプリケーション	<b>適用あり:</b> IHC-P, IP, ICC/IF, WB
種交差性	<b>交差種:</b> Human <b>交差が予測される動物種:</b> Mouse, Rat, Macaque monkey 
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	Recombinant Human hnRNP A1 (isoform A1-A) protein ( <a href="#">ab91691</a> ) can be used as a positive control in WB. This antibody gave a positive signal in HeLa whole cell lysate and Human normal skin tissue section. ICC/IF positive control: U2OS cells
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&amp;As</p>

### 製品の特性

---

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -80°C. Avoid freeze / thaw cycle.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituent: PBS

Batches of this product that have a concentration < 1mg/ml may have BSA added as a stabilising agent. If you would like information about the formulation of a specific lot, please contact our scientific support team who will be happy to help.

精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

## アプリケーション


**The Abpromise guarantee** **Abpromise保証は、次のテスト済みアプリケーションにおけるab4791の使用に適用されます**  
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use a concentration of 5 µg/ml. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
IP		Use at an assay dependent concentration.
ICC/IF		Use at an assay dependent concentration.
WB	★★★★★ (1)	1/500 - 1/1000. Detects a band of approximately 39 kDa (predicted molecular weight: 41 kDa).

## ターゲット情報

機能	Involved in the packaging of pre-mRNA into hnRNP particles, transport of poly(A) mRNA from the nucleus to the cytoplasm and may modulate splice site selection. May play a role in HCV RNA replication.
配列類似性	Contains 2 RRM (RNA recognition motif) domains.
翻訳後修飾	Arg-194, Arg-206 and Arg-225 are dimethylated, probably to asymmetric dimethylarginine. Sumoylated.
細胞内局在	Nucleus. Cytoplasm. Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Shuttles continuously between the nucleus and the cytoplasm along with mRNA. Component of ribonucleosomes. In the course of viral infection, colocalizes with HCV NS5B at speckles in the cytoplasm in a HCV-replication dependent manner.


## 画像



Immunocytochemistry/ Immunofluorescence - Anti-hnRNP A1 antibody (ab4791)

ab4791 staining hnRNP A1 in U2OS cells. The cells were fixed with 4% PFA (10 mins), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab4791 at 5µg/ml and **ab195883**, Rat monoclonal to alpha Tubulin (Alexa Fluor® 488), at 2µg/ml (shown in green). The secondary antibody (shown in red) was **ab150083**, Alexa Fluor® 647 Goat anti-Rabbit IgG (H+L) used at a 1/1000 dilution for 1h at room temperature. Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Western blot - Anti-hnRNP A1 antibody (ab4791)

**All lanes** : Anti-hnRNP A1 antibody (ab4791) at 1/500 dilution

**Lane 1** : HeLa Whole Cell Extract

**Lane 2** : HeLa Nuclear Extract

#### Secondary

**All lanes** : Goat Anti-Rabbit IgG H&L (HRP) (**ab6721**) at 1/2000 dilution

**Predicted band size:** 41 kDa


**Observed band size:** 39 kDa

Rabbit polyclonal to hnRNPA1 (ab4791) at 1/500.

Lane 1: HeLa Whole Cell Extract

Lane 2: HeLa Nuclear Extract

Secondary ab: Goat anti-rabbit IgG HRP conjugate **ab6721** (1/2000)




hnRNP A1 was immunoprecipitated using 0.5mg Hela whole cell extract, 5µg of Rabbit polyclonal to hnRNP A1 and 50µl of protein G magnetic beads (+). No antibody was added to the control (-).

The antibody was incubated under agitation with Protein G beads for 10min, Hela whole cell extract lysate diluted in RIPA buffer was added to each sample and incubated for a further 10min under agitation.

Proteins were eluted by addition of 40µl SDS loading buffer and incubated for 10min at 70°C; 10µl of each sample was separated on a SDS PAGE gel, transferred to a nitrocellulose membrane, blocked with 5% BSA and probed with ab4791.

Secondary: Mouse monoclonal [SB62a] Secondary Antibody to Rabbit IgG light chain (HRP) ([ab99697](#)).

Band: 36kDa: hnRNP A1.



Immunofluorescent imaging of human cells (U2OS) with ab4791 reveals the expected ribonucleoprotein particular staining in the nucleus.


IF was performed with a standard paraformaldehyde technique (fixed in

PBS buffered PFH 4% for 5 minutes, permeabilised with 0.5% triton-PBS

for 5 minutes, blocked with 5% milk / 0.2% tween for one

hour. Primary antibody used at 1/200 in 5% milk / 0.2% TWEEN for one hour, secondary antibody for 30 minutes. All blocking and

incubation steps carried out at 37 degrees. Nuclei counterstained with Hoechst stain (blue).



IHC image of hnRNP A1 staining in human skin FFPE section, performed on a Leica Bond™ system using the standard protocol F. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20 mins. The section was then incubated with ab4791, 5µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-hnRNP A1 antibody (ab4791)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### **Our Abpromise to you: Quality guaranteed and expert technical support**

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
  
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors