

Anti-Nup153 antibody [QE5] ab24700

★★★★★ **5 Abreviews** **57 References** **画像数 6**

製品の概要

製品名	Anti-Nup153 antibody [QE5]
製品の詳細	Mouse monoclonal [QE5] to Nup153
由来種	Mouse
特異性	This antibody could also recognise other NPC polypeptides, p250 and p62, apart from Nup153.
アプリケーション	適用あり: ICC/IF
種交差性	交差種: Mouse, Human
免疫原	Full length protein. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	ICC/IF: U2OS, HepG2, HeLa, Human and Mouse cells. ICC: NIH3T3 cells
特記事項	<p>This antibody clone is manufactured by Abcam. If you require a custom buffer formulation or conjugation for your experiments, please contact orders@abcam.com.</p> <p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C or -80°C. Avoid repeated freeze / thaw cycles.
バッファー	pH: 7.40 Preservative: 0.02% Sodium azide Constituents: PBS, 6.97% L-Arginine
精製度	Protein G purified
ポリ/モノ	モノクローナル
クローン名	QE5
アイソタイプ	IgG1

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab24700の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF	★★★★★ (2)	Use a concentration of 1 µg/ml.

ターゲット情報


機能	Possible DNA-binding subunit of the nuclear pore complex (NPC). The repeat-containing domain may be involved in anchoring components of the pore complex to the pore membrane.
配列類似性	Contains 4 RanBP2-type zinc fingers.
ドメイン	Contains F-X-F-G repeats.
細胞内局在	Nucleus > nuclear pore complex. Located to the terminal ring structure of the nucleoplasmic cage.

画像



ab24700 staining Nup153 in HeLa cells. The cells were fixed with 100% methanol (5 min) then permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab24700 at 0.4 µg/ml concentration and **ab6046** (Rabbit polyclonal to beta Tubulin) at 1/1000 dilution overnight at +4°C followed by a further incubation at room temperature for 1h with a goat secondary antibody to mouse IgG (Alexa Fluor® 488) (**ab150117**) at 2 µg/ml (shown in green) and a goat secondary antibody to rabbit IgG (Alexa Fluor® 594) (**ab150080**) at 2 µg/ml (shown in red). Nuclear DNA was labelled in blue with DAPI.

Image was taken with a confocal microscope (Leica-Microsystems TCS SP8).




Immunocytochemistry/ Immunofluorescence - Anti-Nup153 antibody [QE5] (ab24700)

ab24700 staining Nup153 in NIH3T3 cells. The cells were fixed with 4% paraformaldehyde (10 min), permeabilized with 0.1% PBS-Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated overnight at 4°C with ab24700 at 1µg/ml and **ab6046**, Rabbit polyclonal to beta Tubulin - Loading Control. Cells were then incubated with **ab150117**, Goat polyclonal Secondary Antibody to Mouse IgG H&L (Alexa Fluor® 488) preadsorbed at 1/1000 dilution (shown in green) and **ab150080**, Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (Alexa Fluor® 594) at 1/1000 dilution (shown in pseudocolour red). Nuclear DNA was labelled with DAPI (shown in blue).

Also suitable in cells fixed with 100% methanol (5 min).


Image was acquired with a confocal microscope (Leica-Microsystems TCS SP8) and a single confocal section is shown.



Immunocytochemistry/ Immunofluorescence - Anti-Nup153 antibody [QE5] (ab24700)

This image is courtesy of an anonymous Abreview.

Immunocytochemical analysis of U2OS cells, labeling Nup153 with ab24700. Cells were formaldehyde fixed, permeabilized with NP40, and blocked with 3% BSA for 1 hour at 21°C. Immunostaining with ab24700 diluted 1/100 for 12 hours at 4°C.




Human and mouse cells treated with different concentrations of ab24700. The cells were fixed in 4% formaldehyde and permeabilized in 0.2% Triton x100 for 10 minutes at room temperature, then washed 3 times in PBS. The fixation and permeabilization was carried out in a single step. The image shows clear staining of the nuclear envelope (green). The DNA is stained with DAPI (blue).

- A: HeLa cells, ab24700 used at 4µg/ml
- B: 3T3 cells, ab24700 used at 2µg/ml
- C: 3T3 cells, ab24700 used at 4µg/ml
- D: 3T3 cells, ab24700 used at 6.6µg/ml


Immunocytochemistry/ Immunofluorescence - Anti-Nup153 antibody [QE5] (ab24700)

This image is courtesy of Patrizia Lavia & Maria de Luca, University La Sapienza



Methanol fixed HeLa stained with ab24700. This antibody brilliantly highlights the nuclear membrane (green). The golgi is stained with Giantin (yellow).

Immunocytochemistry/ Immunofluorescence - Anti-Nup153 antibody [QE5] (ab24700)



Human and mouse cells treated with different concentrations of ab24700. The cells were fixed in 100% methanol for 20 minutes at -20°C, then washed once in PBS. The image shows clear staining of the nuclear envelope (green). The DNA is stained with DAPI (blue).

A: HeLa cells, ab24700 used at 4µg/ml

B: 3T3 cells, ab24700 used at 2µg/ml

C: 3T3 cells, ab24700 used at 4µg/ml

D: 3T3 cells, ab24700 used at 6.6µg/ml

Immunocytochemistry/ Immunofluorescence - Anti-Nup153 antibody [QE5] (ab24700)

This image is courtesy of Patrizia Lavia & Maria de Luca, University La Sapienza

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors