abcam

Product datasheet

LY 341495 disodium salt, group II/group III mGlu antagonist ab120818

1 References 画像数 1

製品の概要

製品名 LY 341495 disodium salt, group II/group III mGlu antagonist

製品の詳細 Water-soluble group II/group III mGlu antagonist. Disodium salt of LY 341495 (ab120199).

生理活性の詳細 Water-soluble and very selective nanomolar potent antagonist for group Il mGlu receptors. Also a

relatively potent antagonist for group III mGlu receptors at high nanomolar to low micromolar concentrations. Disodium salt of LY 341495 ($\underline{ab120199}$) which displays K_d values of 1.67 (human mGlu₂), 0.75 (mGlu₃), 31.6 (mGlu₆), 72.7 (mGlu_{7a}) and 14 (human mGlu_{8a} receptors) nM. LY 341495 displays IC₅₀ values of 7.8, 8.2 and 22 μ M for mGlu_{1a}, mGlu_{5a} and mGlu₄ receptors,

respectively. It is readily brain penetrant and active in vivo.

Also available in simple stock solutions (ab144500) - add 1 ml of water to get an exact, ready-to-

use concentration.

精製度 > 98%

CAS 番号 201943-63-7

構造式 H₂N₂CO₂N

製品の特性

体系名 (2S)-2-Amino-2-[(1S,2S)-2-carboxycycloprop-1-yl]-3-(xanth-9-yl)propanoic acid disodium salt

分子量 397.34

分子式 C₂₀H₁₇NNa₂O₅

PubChem 登録番号 9819927

保存方法 Store at -20°C. Store under desiccating conditions. The product can be stored for up to 12

months.

溶解性 Soluble in water to 100 mM

使用に関する注意 Wherever possible, you should prepare and use solutions on the same day. However, if you need

to make up stock solutions in advance, we recommend that you store the solution as aliquots in

tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and prior to opening the vial we recommend that you allow your product to equilibrate to room temperature for at least 1 hour.

Need more advice on solubility, usage and handling? Please visit our <u>frequently asked</u> <u>questions (FAQ) page</u> for more details.


SMILES 線形表記

由来

 $\verb|C1[C@@H]([C@H]1[C@@](CC2C3=CC=CC=C3OC4=CC=CC=C24)(C(=O)O)N)C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C(=O)O|C($

Synthetic

画像



2D chemical structure image of ab120818, LY 341495 disodium salt, group Il/group Ill mGlu antagonist

Chemical Structure - LY 341495 disodium salt, group II/group III mGlu antagonist (ab120818)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES, NOT FOR USE IN HUMANS"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

- · Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to

identify how to use these products effectively. If you require further assistance please contact the scientific support team