

Paclitaxel, Anticancer agent ab120143

★★★★★ [1 Abreviews](#) [14 References](#) [画像数 2](#)


製品の概要

製品名 Paclitaxel, Anticancer agent

製品の詳細 Anticancer agent

CAS 番号 33069-62-4

構造式



製品の特性

体系名 5β,20-Epoxy-1,2a,4,7β,10β,13a-hexahydroxytax-11-en-9-one 4,10-diacetate 2-benzoate 13-ester with (2R,3S)-N-benzoyl-3-phenylisoserine

分子量 853.91

分子式 C₄₇H₅₁NO₁₄

PubChem 登録番号 36314

保存方法 Store at -20°C. Store under desiccating conditions. The product can be stored for up to 12 months.

溶解性 Soluble in DMSO to 100 mM and in ethanol to 25 mM

使用に関する注意 Wherever possible, you should prepare and use solutions on the same day. However, if you need to make up stock solutions in advance, we recommend that you store the solution as aliquots in tightly sealed vials at -20°C. Generally, these will be useable for up to one month. Before use, and prior to opening the vial we recommend that you allow your product to equilibrate to room temperature for at least 1 hour.

Refer to SDS for further information.

Note: Some researchers have found solubility issues in PBS. The recommended procedure (from researcher feedback) for successful solubilisation is as follows: Solubilise 10 mg of ab120143 in 1.25 mL ethanol absolute, mix well, add 1.25mL Cremophor, homogenise. Store at -20°C and after 24 h, defreeze quickly and solubilise in PBS, no crystallisation observed. Please note,

Abcam has not yet tested this method.

Need more advice on solubility, usage and handling? Please visit our [frequently asked questions \(FAQ\) page](#) for more details.


SMILES 線形表記

```
O=C(N[C@@H](c1ccccc1)[C@@H](O)C(=O)O[C@H]5C[C@@]6(O)[C@@H](OC(=O)c2ccccc2)[C@H]3[C@@](C)([C@@H](O)C[C@H]4OC[C@@]34OC(C)=O)C(=O)[C@H](OC(C)=O)C(=C5C)C6(C)C)c7ccccc7
```

由来


Synthetic

画像



2D chemical structure image of ab120143, Paclitaxel, Anticancer agent

Chemical Structure - Paclitaxel, Anticancer agent
(ab120143)



Functional Studies - Paclitaxel, Anticancer agent
(ab120143)

Dong et al PLoS One. 2013; 8(12): e83519.
Reproduced under the Creative Commons license
<http://creativecommons.org/licenses/by/4.0/>

E2 inhibits Paclitaxel induced androgen-independent prostate cancer cell death.

(A-D) 100 nM of E2 was added to the media of (A and B) LNCaP and (C and D) PC3 cells for 96h, followed by addition of **Paclitaxel** at the indicated concentrations for 24h. The cells were stained with Hoechst 33258 (5 µg/ml) to visualize nuclei and propidium iodide (PI) (0.2 µg/ml) to detect membrane damage (B and D). Cell death was quantified by scoring the number of PI positive cells relative to the total number cell nuclei in the same visual field (A and C). The values represent the mean ± S.E. of at least three independent experiments. * denotes p<0.05, ** denotes p<0.01, and *** denotes p<0.001.

E2 = 17-β-estradiol

(From Figure 1 of Dong et al).

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors
- Abcam biochemicals are novel compounds and we have not tested their biological activity in house. Please use the literature to identify how to use these products effectively. If you require further assistance please contact the scientific support team