# abcam

# Product datasheet

# Human AGO3 knockout HeLa cell line ab265320

#### 画像数 2

#### 製品の概要

特記事項

製品名 Human AGO3 knockout HeLa cell line

Parental Cell Line HeLa
Organism Human

Mutation description Knockout achieved by using CRISPR/Cas9, Homozygous: 14 bp deletion in exon 12

Passage number <20

**Knockout validation** Sanger Sequencing, Western Blot (WB)

アプリケーション **適用あり**: WB

Biosafety level

-

**Recommended control:** Human wild-type HeLa cell line (<u>ab255928</u>). Please note a wild-type cell line is not automatically included with a knockout cell line order, if required please add recommended wild-type cell line at no additional cost using the code WILDTYPE-TMTK1.

**Cryopreservation cell medium:** Cell Freezing Medium-DMSO Serum free media, contains 8.7% DMSO in MEM supplemented with methyl cellulose.

Culture medium: DMEM (High Glucose) + 10% FBS

**Initial handling guidelines:** Upon arrival, the vial should be stored in liquid nitrogen vapor phase and not at -80°C. Storage at -80°C may result in loss of viability.

- 1. Thaw the vial in  $37^{\circ}\text{C}$  water bath for approximately 1-2 minutes.
- 2. Transfer the cell suspension (0.8 mL) to a 15 mL/50 mL conical sterile polypropylene centrifuge tube containing 8.4 mL pre-warmed culture medium, wash vial with an additional 0.8 mL culture medium (total volume 10 mL) to collect remaining cells, and centrifuge at 201 x g (rcf) for 5 minutes at room temperature. 10 mL represents minimum recommended dilution. 20 mL represents maximum recommended dilution.
- 3. Resuspend the cell pellet in 5 mL pre-warmed culture medium and count using a haemocytometer or alternative cell counting method. Based on cell count, seed cells in an appropriate cell culture flask at a density of 2x10<sup>4</sup> cells/cm<sup>2</sup>. Seeding density is given as a guide only and should be scaled to align with individual lab schedules.
- 4. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>. Cultures should be monitored daily.

#### Subculture guidelines:

All seeding densities should be based on cell counts gained by established methods. A guide seeding density of  $2x10^4$  cells/cm<sup>2</sup> is recommended.

A partial media change 24 hours prior to subculture may be helpful to encourage growth, if required.

1

Cells should be passaged when they have achieved 80-90% confluence.

This product is subject to limited use licenses from The Broad Institute and ERS Genomics Limited, and is developed with patented technology. For full details of the limited use licenses and relevant patents please refer to our <u>limited use license</u> and <u>patent pages</u>.

We will provide viable cells that proliferate on revival.

#### 製品の特性

Number of cells 1 x 10<sup>6</sup> cells/vial, 1 mL

Adherent /Suspension Adherent

Tissue Cervix
Cell type epithelial

**Disease** Adenocarcinoma

**Gender** Female

**STR Analysis** Amelogenin X D5S818: 11, 12 D13S317: 12, 13.3 D7S820: 8, 12 D16S539: 9, 10 vWA: 16, 18

TH01: 7 TPOX: 8,12 CSF1PO: 9, 10

Mycoplasma free Yes

保存方法 Shipped on Dry Ice. Store in liquid nitrogen.

バッファー Constituents: 8.7% Dimethylsulfoxide, 2% Cellulose, methyl ether

### ターゲット情報

機能 Required for RNA-mediated gene silencing (RNAi). Binds to short RNAs such as microRNAs

(miRNAs) and represses the translation of mRNAs which are complementary to them. Lacks

endonuclease activity and does not appear to cleave target mRNAs.

**配列類似性** Belongs to the argonaute family. Ago subfamily.

Contains 1 PAZ domain. Contains 1 Piwi domain.


**細胞内局在** Cytoplasm > P-body.

## アプリケーション

**The Abpromise guarantee** <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab265320の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration. Predicted molecular weight: 97 kDa.

#### 画像



Western blot - Human AGO3 knockout HeLa cell line (ab265320)

**Lanes 1-2 & 4**: Anti-AGO3 antibody [EPR9576] (<u>ab154844</u>) at 1/2000 dilution

Lane 3: 154844 at 1/2000 dilution

Lane 1: Wild-type HeLa lysate

Lane 2 : Argonaute RISC Catalytic Component 3 knockout HeLa

lysate

Lane 3 : Daudi lysate
Lane 4 : A549 lysate

Lysates/proteins at 40 µg per lane.

Performed under reducing conditions.

Predicted band size: 97 kDa

**Lanes 1-4:** Merged signal (red and green). Green - <u>ab154844</u> observed at 97 kDa. Red - loading control <u>ab8245</u> observed at 37 kDa.

<u>ab154844</u> Anti-AGO3 antibody [EPR9576] was shown to specifically react with Argonaute RISC Catalytic Component 3 in wild-type HeLa cells. Loss of signal was observed when knockout cell line ab265320 (knockout cell lysate <u>ab257819</u>) was used. Wild-type and Argonaute RISC Catalytic Component 3 knockout samples were subjected to SDS-PAGE. <u>ab154844</u> and Anti-GAPDH antibody [6C5] - Loading Control (<u>ab8245</u>) were incubated overnight at 4°C at 1 in 2000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit lgG H&L (IRDye® 800CW) preadsorbed (<u>ab216773</u>) and Goat anti-Mouse lgG H&L (IRDye® 680RD) preadsorbed (<u>ab216776</u>) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

Sanger Sequencing - Human AGO3 knockout HeLa cell line (ab265320)

Homozygous: 14 bp deletion in exon 12.

# Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

# Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors