

# Human PTGES3 (p23) knockout HEK-293T cell line ab266791

画像数 4

## 製品の概要

|                      |   |
|----------------------|---|
| 製品名                  | Human PTGES3 (p23) knockout HEK-293T cell line  |
| Parental Cell Line   | HEK293T   |
| Organism             | Human   |
| Mutation description | Knockout achieved by using CRISPR/Cas9, 10 bp insertion in exon 2 and 1 bp insertion in exon 2  |
| Passage number       | <20   |
| Knockout validation  | Sanger Sequencing, Western Blot (WB)  |
| アプリケーション             | 適用あり: WB  |
| Biosafety level      | 2   |
| 特記事項                 | <p><b>Recommended control:</b> Human wild-type HEK293T cell line (<a href="#">ab255449</a>). Please note a wild-type cell line is not automatically included with a knockout cell line order, if required please add recommended wild-type cell line at no additional cost using the code WILDTYPE-TMTK1.</p> <p><b>Cryopreservation cell medium:</b> Cell Freezing Medium-DMSO Serum free media, contains 8.7% DMSO in MEM supplemented with methyl cellulose.</p> <p><b>Culture medium:</b> DMEM (High Glucose) + 10% FBS</p> <p><b>Initial handling guidelines:</b> Upon arrival, the vial should be stored in liquid nitrogen vapor phase and not at -80°C. Storage at -80°C may result in loss of viability.</p> <ol style="list-style-type: none"> <li>1. Thaw the vial in 37°C water bath for approximately 1-2 minutes.</li> <li>2. Transfer the cell suspension (0.8 mL) to a 15 mL/50 mL conical sterile polypropylene centrifuge tube containing 8.4 mL pre-warmed culture medium, wash vial with an additional 0.8 mL culture medium (total volume 10 mL) to collect remaining cells, and centrifuge at 201 x g (rcf) for 5 minutes at room temperature. 10 mL represents minimum recommended dilution. 20 mL represents maximum recommended dilution.</li> <li>3. Resuspend the cell pellet in 5 mL pre-warmed culture medium and count using a haemocytometer or alternative cell counting method. Based on cell count, seed cells in an appropriate cell culture flask at a density of <math>2 \times 10^4</math> cells/cm<sup>2</sup>. Seeding density is given as a guide only and should be scaled to align with individual lab schedules.</li> <li>4. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>. Cultures should be monitored daily.</li> </ol> <p><b>Subculture guidelines:</b></p> <p>All seeding densities should be based on cell counts gained by established methods. A guide seeding density of <math>2 \times 10^4</math> cells/cm<sup>2</sup> is recommended.</p> <p>A partial media change 24 hours prior to subculture may be helpful to encourage growth, if required.</p> |

Cells should be passaged when they have achieved 80-90% confluence.

This product is subject to limited use licenses from The Broad Institute, ERS Genomics Limited and Sigma-Aldrich Co. LLC, and is developed with patented technology. For full details of the licenses and patents please refer to our [limited use license](#) and [patent pages](#).

We will provide viable cells that proliferate on revival.

## 製品の特性

|                      |  |
|----------------------|--|
| Number of cells      | 1 x 10 <sup>6</sup> cells/vial, 1 mL   |
| Adherent /Suspension | Adherent   |
| Tissue               | Kidney   |
| Cell type            | epithelial   |
| STR Analysis         | Amelogenin X D5S818: 8, 9 D13S317: 12, 14 D7S820: 11 D16S539: 9, 13 vWA: 16, 19 TH01: 7, 9.3 TPOX: 11 CSF1PO: 11, 12 |
| Mycoplasma free      | Yes  |
| 保存方法                 | Shipped on Dry Ice. Store in liquid nitrogen.  |
| バッファー                | Constituents: 8.7% Dimethylsulfoxide, 2% Cellulose, methyl ether   |

## ターゲット情報


|       |  |
|-------|--|
| 機能    | Molecular chaperone that localizes to genomic response elements in a hormone-dependent manner and disrupts receptor-mediated transcriptional activation, by promoting disassembly of transcriptional regulatory complexes. |
| パスウェイ | Lipid metabolism; prostaglandin biosynthesis.  |
| 配列類似性 | Belongs to the p23/wos2 family.<br>Contains 1 CS domain.   |
| 細胞内局在 | Cytoplasm.   |

## アプリケーション

**The Abpromise guarantee**      **Abpromise保証は、** 次のテスト済みアプリケーションにおけるab266791の使用に適用されます  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

| アプリケーション | Abreviews | 特記事項   |
|----------|-----------|--|
| WB       |           | Use at an assay dependent concentration. Predicted molecular weight: 19 kDa. |

## 画像



Western blot - Human PTGES3 (p23) knockout  
HEK293T cell line (ab266791)

**All lanes :** Anti-p23 antibody [EPR3846] ([ab92503](#)) at 1/1000 dilution

**Lane 1 :** Wild-type HEK-293T (Human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate

**Lane 2 :** PTGES3 knockout HEK-293T (Human epithelial cell line from embryonic kidney transformed with large T antigen) whole cell lysate

**Lane 3 :** Jurkat (Human T cell leukemia cell line from peripheral blood) whole cell lysate

**Lane 4 :** HeLa (Human epithelial cell line from cervix adenocarcinoma) whole cell lysate

Lysates/proteins at 20 µg per lane.

### Secondary

**All lanes :** Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) at 1/10000 dilution

**Predicted band size:** 19 kDa

**Observed band size:** 23 kDa

**Lanes 1-4:** Merged signal (red and green). Green - [ab92503](#) observed at 23 kDa. Red - loading control [ab8245](#) observed at 36 kDa.

[ab92503](#) Anti-p23 antibody [EPR3846] was shown to specifically react with p23 in wild-type HEK-293T cells. Loss of signal was observed when knockout cell line ab266791 (knockout cell lysate [ab258151](#)) was used. Wild-type and p23 knockout samples were subjected to SDS-PAGE. [ab92503](#) and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

|     |  |
|-----|--|
| Mut | TTATCCTTTTGTAGGCAGCCTGCTTCTGCAAAAGTGGTACGATCGAAGGGACTATGTCTT |
| WT  | TTATCCTTTTGTAGGCAGCCTGCTTCTGCAAA GTGGTACGATCGAAGGGACTATGTCTT |


Sanger Sequencing - Human PTGES3 knockout  
HEK293T cell line (ab266791)

Allele-1: 1 bp insertion in exon 2

|     |   |
|-----|---|
| Mut | TTATCCTTTTGTAGGCAGCCTGCTTCTGCAAAAGTGGTACAAAGTGGTACGATCGAAGGGA |
| WT  | TTATCCTTTTGTAGGCAGCCTGCTTCTGCAAA GTGGTACGATCGAAGGGA           |

Sanger Sequencing - Human PTGES3 knockout  
HEK293T cell line (ab266791)

Allele-2: 10 bp insertion in exon 2.



**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

## Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise,

please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

#### **Terms and conditions**

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors