

# Human TGM2 (Transglutaminase 2) knockout A549 cell line ab267110

画像数 4

### 製品の概要

製品名	Human TGM2 (Transglutaminase 2) knockout A549 cell line
Parental Cell Line	A549
Organism	Human
Mutation description	Knockout achieved by using CRISPR/Cas9, 1 bp insertion in exon6.
Passage number	<20
Knockout validation	Sanger Sequencing, Western Blot (WB)
アプリケーション	適用あり: WB
Biosafety level	1
特記事項	<p><b>Recommended control:</b> Human wild-type A549 cell line (<a href="#">ab255450</a>). Please note a wild-type cell line is not automatically included with a knockout cell line order, if required please add recommended wild-type cell line at no additional cost using the code WILDTYPE-TMTK1.</p> <p><b>Cryopreservation cell medium:</b> Cell Freezing Medium-DMSO Serum free media, contains 8.7% DMSO in MEM supplemented with methyl cellulose.</p> <p><b>Culture medium:</b> F-12K + 10% FBS</p> <p><b>Initial handling guidelines:</b> Upon arrival, the vial should be stored in liquid nitrogen vapor phase and not at -80°C. Storage at -80°C may result in loss of viability.</p> <ol style="list-style-type: none"><li>1. Thaw the vial in 37°C water bath for approximately 1-2 minutes.</li><li>2. Transfer the cell suspension (0.8 mL) to a 15 mL/50 mL conical sterile polypropylene centrifuge tube containing 8.4 mL pre-warmed culture medium, wash vial with an additional 0.8 mL culture medium (total volume 10 mL) to collect remaining cells, and centrifuge at 201 x g (rcf) for 5 minutes at room temperature. 10 mL represents minimum recommended dilution. 20 mL represents maximum recommended dilution.</li><li>3. Resuspend the cell pellet in 5 mL pre-warmed culture medium and count using a haemocytometer or alternative cell counting method. Based on cell count, seed cells in an appropriate cell culture flask at a density of <math>2 \times 10^3</math>-<math>1 \times 10^4</math> cells/cm<sup>2</sup>. Seeding density is given as a guide only and should be scaled to align with individual lab schedules.</li><li>4. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>. Cultures should be monitored daily.</li></ol> <p><b>Subculture guidelines:</b></p> <p>All seeding densities should be based on cell counts gained by established methods. A guide seeding density of <math>6 \times 10^4</math> cells/cm<sup>2</sup> is recommended.</p>

A partial media change 24 hours prior to subculture may be helpful to encourage growth, if required.

Cells should be passaged when they have achieved 80-90% confluence.

Do not exceed  $7 \times 10^4$  cells/cm<sup>2</sup>.

This product is subject to limited use licenses from The Broad Institute, ERS Genomics Limited and Sigma-Aldrich Co. LLC, and is developed with patented technology. For full details of the licenses and patents please refer to our [limited use license](#) and [patent pages](#).

We will provide viable cells that proliferate on revival.

## 製品の特性

Number of cells	1 x 10 <sup>6</sup> cells/vial, 1 mL
Adherent /Suspension	Adherent
Tissue	Lung
Cell type	epithelial
Disease	Carcinoma
Gender	Male
STR Analysis	Amelogenin X,YD5S818: 11 D13S317: 11 D7S820: 8, 11 D16S539: 11, 12 WWA: 14 TH01: 8,9.3 TPOX: 8,11 CSF1PO: 10, 12
Antibiotic resistance	Puromycin 1.00µg/ml
Mycoplasma free	Yes
保存方法	Shipped on Dry Ice. Store in liquid nitrogen.
バッファー	Constituents: 8.7% Dimethylsulfoxide, 2% Cellulose, methyl ether

## ターゲット情報


機能	Catalyzes the cross-linking of proteins and the conjugation of polyamines to proteins.
配列類似性	Belongs to the transglutaminase superfamily. Transglutaminase family.

## アプリケーション

**The Abpromise guarantee**      **Abpromise保証は、次のテスト済みアプリケーションにおけるab267110の使用に適用されます**  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration. Predicted molecular weight: 77 kDa.

## 画像



Western blot - Human TGM2 knockout A549 cell line (ab267110)

**All lanes** : Anti-Transglutaminase 2 antibody [EP2957] (**ab109200**) at 1/10000 dilution

**Lane 1** : Wild-type A549 cell lysate

**Lane 2** : TGM2 knockout A549 cell lysate

Lysates/proteins at 20 µg per lane.


Performed under reducing conditions.

**Predicted band size:** 77 kDa

**Observed band size:** 77 kDa

**Lanes 1-2:** Merged signal (red and green). Green - **ab109200** observed at 77 kDa. Red - loading control **ab8245** observed at 37 kDa.

**ab109200** Anti-Transglutaminase 2 antibody [EP2957] was shown to specifically react with Transglutaminase 2 in wild-type A549 cells. Loss of signal was observed when knockout cell line ab267110 (knockout cell lysate **ab257087**) was used. Wild-type and Transglutaminase 2 knockout samples were subjected to SDS-PAGE. **ab109200** and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) were incubated overnight at 4°C at 1 in 10000 and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Western blot - Human TGM2 knockout A549 cell line (ab267110)

**All lanes** : Anti-Transglutaminase 2 antibody [EPR2956] (**ab109121**) at 1/1000 dilution

**Lane 1** : Wild-type A549 cell lysate

**Lane 2** : TGM2 knockout A549 cell lysate

Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

**Predicted band size:** 77 kDa

**Observed band size:** 77 kDa

**Lanes 1-2:** Merged signal (red and green). Green - **ab109121** observed at 77 kDa. Red - loading control **ab8245** observed at 37 kDa.


**ab109121** Anti-Transglutaminase 2 antibody [EPR2956] was shown to specifically react with Transglutaminase 2 in wild-type A549 cells. Loss of signal was observed when knockout cell line ab267110 (knockout cell lysate **ab257087**) was used. Wild-type and Transglutaminase 2 knockout samples were subjected to SDS-PAGE. **ab109121** and Anti-GAPDH antibody [6C5] - Loading Control (**ab8245**) were incubated overnight at 4°C at 1 in 1000 and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed (**ab216773**) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed (**ab216776**) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.

```

Mut  GTGTGCTGCTGGGACGCTGGGACAACAACCTACGGGGACGGCGTCAGCCCCATGTCCTGG
      |||
WT   GTGTGCTGCTGGGACGCTGGGACAACAACCTACGGGGACGGCGTCAGCCCCATGTCCTGG
  
```

Sanger Sequencing - Human TGM2 knockout A549 cell line (ab267110)

Homozygous: 1 bp insertion in exon6



Representative images of TGM2 knockout A549 cells, low and high confluency examples (top left and right respectively) and wild-type A549 cells, low and high confluency (bottom left and right respectively) showing typical adherent, epithelial-like morphology. Images were captured at 10X magnification using a EVOS XL Core microscope.

Cell Culture - Human TGM2 (Transglutaminase 2)  
knockout A549 cell line (ab267110)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors