# abcam

# Product datasheet

# Human TNFRSF10B (DR5) knockout HeLa cell line ab264922

### 画像数 2

#### 製品の概要

製品名 Human TNFRSF10B (DR5) knockout HeLa cell line

Parental Cell Line HeLa
Organism Human

Mutation description Knockout achieved by using CRISPR/Cas9, Homozygous: Insertion of the selection cassette in

exon 1

Passage number <20

**Knockout validation** Sanger Sequencing, Western Blot (WB)

アプリケーション **適用あり**: WB

Biosafety level 2

特記事項

**Recommended control:** Human wild-type HeLa cell line (<u>ab255448</u>). Please note a wild-type cell line is not automatically included with a knockout cell line order, if required please add recommended wild-type cell line at no additional cost using the code WILDTYPE-TMTK1.

**Cryopreservation cell medium:** Cell Freezing Medium-DMSO Serum free media, contains 8.7% DMSO in MEM supplemented with methyl cellulose.

Culture medium: DMEM (High Glucose) + 10% FBS

**Initial handling guidelines:** Upon arrival, the vial should be stored in liquid nitrogen vapor phase and not at -80°C. Storage at -80°C may result in loss of viability.

- 1. Thaw the vial in 37°C water bath for approximately 1-2 minutes.
- 2. Transfer the cell suspension (0.8 mL) to a 15 mL/50 mL conical sterile polypropylene centrifuge tube containing 8.4 mL pre-warmed culture medium, wash vial with an additional 0.8 mL culture medium (total volume 10 mL) to collect remaining cells, and centrifuge at 201 x g (rcf) for 5 minutes at room temperature. 10 mL represents minimum recommended dilution. 20 mL represents maximum recommended dilution.
- 3. Resuspend the cell pellet in 5 mL pre-warmed culture medium and count using a haemocytometer or alternative cell counting method. Based on cell count, seed cells in an appropriate cell culture flask at a density of 2x10<sup>4</sup> cells/cm<sup>2</sup>. Seeding density is given as a guide only and should be scaled to align with individual lab schedules.
- 4. Incubate the culture at 37°C incubator with 5% CO<sub>2</sub>. Cultures should be monitored daily.

## Subculture guidelines:

All seeding densities should be based on cell counts gained by established methods. A guide seeding density of  $2x10^4$  cells/cm<sup>2</sup> is recommended.

A partial media change 24 hours prior to subculture may be helpful to encourage growth, if

1

required.

Cells should be passaged when they have achieved 80-90% confluence.

This product is subject to limited use licenses from The Broad Institute, ERS Genomics Limited and Sigma-Aldrich Co. LLC, and is developed with patented technology. For full details of the licenses and patents please refer to our <u>limited use license</u> and <u>patent pages</u>.

We will provide viable cells that proliferate on revival.

### 製品の特性

Number of cells 1 x 10<sup>6</sup> cells/vial, 1 mL

Adherent /Suspension Adherent
Tissue Cervix
Cell type epithelial

**Disease** Adenocarcinoma

**Gender** Female

**STR Analysis** Amelogenin X D5S818: 11, 12 D13S317: 12, 13.3 D7S820: 8, 12 D16S539: 9, 10 vWA: 16, 18

TH01: 7 TPOX: 8, 12 CSF1PO: 9, 10

Mycoplasma free Yes

保存方法 Shipped on Dry Ice. Store in liquid nitrogen.

パップァー Constituents: 8.7% Dimethylsulfoxide, 2% Cellulose, methyl ether

#### ターゲット情報

機能 Receptor for the cytotoxic ligand TNFSF10/TRAIL. The adapter molecule FADD recruits

caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-

kappa-B. Essential for ER stress-induced apoptosis.

組織特異性 Widely expressed in adult and fetal tissues; very highly expressed in tumor cell lines such as

HeLaS3, K-562, HL-60, SW480, A-549 and G-361; highly expressed in heart, peripheral blood lymphocytes, liver, pancreas, spleen, thymus, prostate, ovary, uterus, placenta, testis, esophagus,

stomach and throughout the intestinal tract; not detectable in brain.

**関連疾患** Squamous cell carcinoma of the head and neck

配列類似性 Contains 1 death domain.

Contains 3 TNFR-Cys repeats.


細胞内局在 Membrane.

#### アプリケーション

**The Abpromise guarantee** <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab264922の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use at an assay dependent concentration. Predicted molecular weight: 47 kDa.

#### 画像



Western blot - Human TNFRSF10B (DR5) knockout HeLa cell line (ab264922) **All lanes :** Anti-DR5 antibody [EPR19310] (**ab199357**) at 1/1000 dilution

Lane 1 : Wild-type HeLa cell lysate

Lane 2 : DR5 knockout HeLa cell lysate


Lysates/proteins at 20 µg per lane.

Performed under reducing conditions.

**Predicted band size:** 47 kDa **Observed band size:** 47 kDa

**Lanes 1-2:** Merged signal (red and green). Green - <u>ab199357</u> observed at 47 kDa. Red - Anti-GAPDH antibody [6C5] - Loading Control (<u>ab8245</u>) observed at 37 kDa.

ab199357 was shown to react with DR5 in wild-type HeLa cells in western blot. Loss of signal was observed when knockout cell line ab264922 (knockout cell lysate ab257748) was used. Wild-type HeLa and TNFRSF10B knockout HeLa cell lysates were subjected to SDS-PAGE. Membrane was blocked for 1 hour at room temperature in 0.1% TBST with 3% non-fat dried milk. ab199357 and Anti-GAPDH antibody [6C5] - Loading Control (ab8245) overnight at 4°C at a 1 in 1000 dilution and a 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye®800CW) preadsorbed (ab216773) and Goat anti-Mouse IgG H&L (IRDye®680RD) preadsorbed (ab216776) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Homozygous: Insertion of the selection cassette in exon 1.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

# Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- · Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

#### Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors