

Rad51 overexpression 293T lysate (whole cell) ab94044

画像数 2

医薬用外劇物

製品の概要

| | |
|----------|--|
| 製品名 | Rad51 overexpression 293T lysate (whole cell) |
| 特記事項 | ab94044 is a 293T cell transfected lysate in which Human Rad51 has been transiently over-expressed using a pCMV-Rad51 plasmid. The lysate is provided in 1X Sample Buffer. |
| アプリケーション | 適用あり: WB |

製品の特性


| | |
|-----------------|---|
| Mycoplasma free | Yes |
| 製品の状態 | Liquid |
| 保存方法 | Shipped on dry ice. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles. |
| バッファー | Constituents: 0.01% Bromophenol blue, 2.3% Beta mercaptoethanol, 2% Sodium lauryl sulfate, 0.788% Tris HCl, 10% Glycerol (glycerin, glycerine) |
| 背景 | Disease: Breast cancer Mirror movements 2 Defects in RAD51 are found in a patient with microcephaly, mental retardation without bone marrow failure and pediatric cancers. Domain: The nuclear localization may reside in the C-terminus (between 259 and 339 AA). Function: Plays an important role in homologous strand exchange, a key step in DNA repair through homologous recombination. Binds to single and double-stranded DNA and exhibits DNA-dependent ATPase activity. Catalyzes the recognition of homology and strand exchange between homologous DNA partners to form a joint molecule between a processed DNA break and the repair template. Binds to single-stranded DNA in an ATP-dependent manner to form nucleoprotein filaments which are essential for the homology search and strand exchange (PubMed:26681308). Part of a PALB2-scaffolded HR complex containing BRCA2 and RAD51C and which is thought to play a role in DNA repair by HR. Plays a role in regulating mitochondrial DNA copy number under conditions of oxidative stress in the presence of RAD51C and XRCC3. PTM: Ubiquitinated by the SCF(FBXO18) E3 ubiquitin ligase complex, regulating RAD51 subcellular location and preventing its association with DNA. Phosphorylated. Phosphorylation of Thr-309 by CHEK1 may enhance association with chromatin at sites of DNA damage and promote DNA repair by homologous recombination. Phosphorylation by ABL1 inhibits function. Similarity: Belongs to the RecA family. RAD51 subfamily. Contains 1 HhH domain. Tissue specificity: Highly expressed in testis and thymus, followed by small intestine, placenta, colon, pancreas and ovary. Weakly expressed in breast. |

アプリケーション

The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab94044の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。


| アプリケーション | Abreviews | 特記事項 |
|----------|-----------|-------------------------------------|
| WB | | Use at an assay dependent dilution. |

画像



SDS-PAGE - Rad51 overexpression 293T lysate
(whole cell) (ab94044)

ab94044 at 15µg/lane on an SDS-PAGE gel.



Western blot - Rad51 overexpression 293T lysate
(whole cell) (ab94044)

All lanes : Anti-Rad51 antibody - BSA and Azide free (**ab88572**) at 1/500 dilution

Lane 1 : ab94044

Lane 2 : non transfected lysate

Lysates/proteins at 25 µg per lane.

Secondary

All lanes : non-Abcam, Goat Anti-mouse IgG (H and L) HRP conjugated at 1/2500 dilution

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you

- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors