


DRAQ5™ ab108410

★★★★★ [5 Abreviews](#) [53 References](#) [画像数 3](#)

製品の概要

製品名	DRAQ5™
アプリケーション	適用あり: FM, Flow Cyt, ICC/IF
特記事項	<p>DRAQ5™ is a cell permeable far-red fluorescent DNA dye that can be used in fixed or non-fixed/ live cells in combination with common labels such as GFP or FITC.</p> <p>As with any cell-permeant DNA intercalating probe, DRAQ5 may inhibit cell division in long-term assays and should be tested for any effect.</p> <p>DRAQ5 staining can be used in flow cytometry, live cell imaging and cell-based assays and the dye is highly compatible with standard protocols across many instrumentation platforms.</p> <p>The chemical name of DRAQ5 is <i>1, 5-bis[[2-(di-methylamino)ethyl]amino]-4, 8-dihydroxyanthracene-9, 10-dione</i>.</p> <p>The advantages of DRAQ5 staining include</p> <ul style="list-style-type: none"> - convenient ready-to-use aqueous solution - rapid uptake into living cells, providing a high level of nuclear discrimination - no photobleaching effect - can be used in most cell types, eukaryotic and prokaryotic: mammalian, bacterial, parasitic, plant, etc. - no compensation needed with common FITC/GFP + PE combinations in flow cytometry - no RNase treatment required - no fluorescence enhancement upon DNA binding - compatible with optics of benchtop flow, laser scanning cytometers and non-UV laser scanning and lamp-based confocal microscopes <p>SPECTRAL PROPERTIES:</p> <p>Excitation</p> <p>647 nm line optimal (Ex_{max} 646 nm)</p> <p>488, 514, 568 and 633 nm lines, sub-optimal</p> <p>Two-photon excitation (1047 nm) and excitation dark (700-850 nm)</p> <p>Emission (instrument dependent):</p> <ul style="list-style-type: none"> - 665 nm to infra-red max 681 nm / 697 nm intercalated with dsDNA) - minimal overlap with vis range e.g. GFP and FITC - Em. filters may include 695L, 715LP or 780 LP



Concentration: 5 mM

製品の特徴

製品の状態

Liquid

保存方法

Store at +4°C. Do Not Freeze. Store In the Dark.

アプリケーション


The Abpromise guarantee

Abpromise保証は、次のテスト済みアプリケーションにおけるab108410の使用に適用されます

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。


アプリケーション	Abreviews	特記事項
FM		Use at an assay dependent concentration.
Flow Cyt		1/250 - 1/1000. For Gating of Nucleated cells = 5µM; For Cell Cycle Analysis = 20µM
ICC/IF		1/1000. For Cell-based assays, Immunofluorescence microscopy and In-Cell WB = 5µM It is highly recommended that the concentration and labelling conditions are carefully determined by each investigator for optimal performance in the assay of interest. For more specific information about the applications, please refer to the Protocol Booklet section.

画像




Immunocytochemistry/ Immunofluorescence -
DRAQ5™ (ab108410)

HeLa cells were stained with **Lamin B1 antibody - Nuclear Envelope Marker (ab16048)** and **alpha Tubulin antibody [DM1A] - Loading Control (ab7291)**. The cells were 100% methanol fixed (5 min) and then incubated in 1% BSA in 0.1% PBS-Tween for 1h to permeabilize the cells and block non-specific protein-protein interactions. The cells were then incubated with the primary antibodies (**ab16048** & **ab7291**) at 1µg/ml overnight at 4C. The secondary antibodies were **Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (DyLight® 488), pre-adsorbed (ab96899)** (green) and **Goat polyclonal Secondary Antibody to Mouse IgG - H&L (DyLight® 594), pre-adsorbed (ab96881)** (red) used at 1/250 dilution for 1h at room temperature. 5µM DRAQ5 was added to the secondary antibody mixture to label nuclear DNA (pseudocolor purple).



Immunocytochemistry/ Immunofluorescence -
DRAQ5™ (ab108410)

HeLa cells were stained with **beta Catenin antibody (ab16051)** and **alpha Tubulin antibody [DM1A] - Loading Control (ab7291)**. The cells were 100% methanol fixed (5 min) and then incubated in 1% BSA in 0.1% PBS-Tween for 1h to permeabilize the cells and block non-specific protein-protein interactions. The cells were then incubated with the primary antibodies (**ab16051** & **ab7291**) at 1µg/ml overnight at 4C. The secondary antibodies were **Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (DyLight® 488), pre-adsorbed (ab96899)** (green) and **Goat polyclonal Secondary Antibody to Rabbit IgG - H&L (DyLight® 594), pre-adsorbed (ab96899)** (red) used at 1/250 dilution for 1h at room temperature. 5µM DRAQ5 was added to the secondary antibody mixture to label nuclear DNA (pseudocolor orange).



DRAQ5™-stained nuclei in a adult *Drosophila* brain.

Immunocytochemistry/ Immunofluorescence -
DRAQ5™ (ab108410)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors