

Anti-58K Golgi protein antibody [EPR7909] - BSA and Azide free ab248255

リコンビナント RabMAb

画像数 5

製品の概要

製品名	Anti-58K Golgi protein antibody [EPR7909] - BSA and Azide free
製品の詳細	Rabbit monoclonal [EPR7909] to 58K Golgi protein - BSA and Azide free
由来種	Rabbit
特異性	This antibody is not suitable for testing cell line samples in western blot.
アプリケーション	適用あり: IHC-P, WB 適用なし: Flow Cyt or IP
種交差性	交差種: Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
特記事項	<p>ab248255 is the carrier-free version of ab129005.</p> <p>Our carrier-free antibodies are typically supplied in a PBS-only formulation, purified and free of BSA, sodium azide and glycerol. The carrier-free buffer and high concentration allow for increased conjugation efficiency.</p> <p>This conjugation-ready format is designed for use with fluorochromes, metal isotopes, oligonucleotides, and enzymes, which makes them ideal for antibody labelling, functional and cell-based assays, flow-based assays (e.g. mass cytometry) and Multiplex Imaging applications.</p> <p>Use our conjugation kits for antibody conjugates that are ready-to-use in as little as 20 minutes with <1 minute hands-on-time and 100% antibody recovery: available for fluorescent dyes, HRP, biotin and gold.</p> <p>This product is compatible with the Maxpar[®] Antibody Labeling Kit from Fluidigm, without the need for antibody preparation. Maxpar[®] is a trademark of Fluidigm Canada Inc.</p> <p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C. Do Not Freeze.
バッファー	pH: 7.2 Constituent: PBS
キャリア・フリー	はい
精製度	Protein A purified
ポリクローナル	モノクローナル
クローン名	EPR7909
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab248255の使用に適用されず**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.
WB		Use at an assay dependent concentration. Detects a band of approximately 58 kDa (predicted molecular weight: 58 kDa).

追加情報 Is unsuitable for Flow Cyt or IP.

ターゲット情報

機能	Folate-dependent enzyme, that displays both transferase and deaminase activity. Serves to channel one-carbon units from formiminoglutamate to the folate pool. Binds and promotes bundling of vimentin filaments originating from the Golgi.
パスウェイ	Amino-acid degradation; L-histidine degradation into L-glutamate; L-glutamate from N-formimidoyl-L-glutamate (transferase route): step 1/1. One-carbon metabolism; tetrahydrofolate interconversion.
関連疾患	Defects in FTCD are the cause of glutamate formiminotransferase deficiency (FIGLU-URIA) [MIM:229100]; also known as formiminoglutamicaciduria (FIGLU-uria). It is an autosomal recessive disorder. Features of a severe phenotype, include elevated levels of formiminoglutamate (FIGLU) in the urine in response to histidine administration, megaloblastic anemia, and mental retardation. Features of a mild phenotype include high urinary excretion of FIGLU in the absence of histidine administration, mild developmental delay, and no hematological abnormalities.


配列類似性

In the C-terminal section; belongs to the cyclodeaminase/cyclohydrolase family.
In the N-terminal section; belongs to the formiminotransferase family.

細胞内局在

Cytoplasm > cytoskeleton > centrosome > centriole. Golgi apparatus. More abundantly located around the mother centriole.

画像



Western blot - Anti-58K Golgi protein antibody [EPR7909] - BSA and Azide free (ab248255)

All lanes : Anti-58K Golgi protein antibody [EPR7909] (**ab129005**) at 1/1000 dilution

Lane 1 : Human liver tissue lysate

Lane 2 : Human kidney tissue lysate

Lysates/proteins at 20 µg per lane.

Secondary


All lanes : Goat Anti-Rabbit IgG (HRP) with minimal cross-reactivity with human IgG at 1/2000 dilution

Predicted band size: 58 kDa

Observed band size: 58 kDa

Exposure time: 20 seconds

Blocking buffer and dilution buffer concentration: 5% NFDm/TBST.



Western blot - Anti-58K Golgi protein antibody [EPR7909] - BSA and Azide free (ab248255)

Anti-58K Golgi protein antibody [EPR7909] (**ab129005**) at 1/1000 dilution + Human fetal liver tissue lysate at 10 µg


Secondary

Goat anti-Rabbit HRP at 1/2000 dilution

Predicted band size: 58 kDa


Observed band size: 58 kDa

This data was developed using **ab129005**, the same antibody clone in a different buffer formulation.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-58K Golgi protein antibody [EPR7909] - BSA and Azide free (ab248255)

This data was developed using **ab129005**, the same antibody clone in a different buffer formulation. **ab129005**, at 1/250 dilution, staining 58K Golgi protein in paraffin-embedded Human kidney tissue by Immunohistochemistry. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-58K Golgi protein antibody [EPR7909] - BSA and Azide free (ab248255)

This data was developed using **ab129005**, the same antibody clone in a different buffer formulation. **ab129005**, at 1/250 dilution, staining 58K Golgi protein in paraffin-embedded Human liver tissue by Immunohistochemistry. Perform heat mediated antigen retrieval before commencing with IHC staining protocol.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-58K Golgi protein antibody [EPR7909] - BSA and Azide free (ab248255)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors