



Anti-ABCB4 antibody [P3II-26] ab24108

6 References [画像数 1](#)

製品の概要

製品名	Anti-ABCB4 antibody [P3II-26]
製品の詳細	Mouse monoclonal [P3II-26] to ABCB4
由来種	Mouse
特異性	Clone P3II-26 does not cross-react with the human ABCB1.
アプリケーション	適用あり: Flow Cyt 適用なし: IHC-P
種交差性	交差種: Human
免疫原	Recombinant fragment within Human ABCB4 aa 600-700 (internal sequence). The exact immunogen sequence used to generate this antibody is proprietary information. If additional detail on the immunogen is needed to determine the suitability of the antibody for your needs, please <u>contact</u> our Scientific Support team to discuss your requirements.  <u>Run BLAST with</u>  <u>Run BLAST with</u>
エピトープ	Clone P3II-26 reacts with an internal epitope of ABCB4.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term.
バッファー	pH: 7.3 Preservative: 0.1% Sodium azide Constituent: 0.7% BSA Serum free tissue culture supernatant

精製度	Protein G purified
ポリ/モノ	モノクローナル
クローン名	P3II-26
アイソタイプ	IgG2b

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab24108の使用に適用されます**

アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
Flow Cyt		Use 1µg for 10 ⁶ cells. ab170192 - Mouse monoclonal IgG2b, is suitable for use as an isotype control with this antibody.

追加情報 Is unsuitable for IHC-P.

ターゲット情報

機能	Mediates ATP-dependent export of organic anions and drugs from the cytoplasm. Hydrolyzes ATP with low efficiency. Human MDR3 is not capable of conferring drug resistance. Mediates the translocation of phosphatidylcholine across the canalicular membrane of the hepatocyte.
関連疾患	Defects in ABCB4 are the cause of progressive familial intrahepatic cholestasis type 3 (PFIC3) [MIM:602347]. PFIC3 is an autosomal recessive liver disorder presenting with early onset cholestasis that progresses to cirrhosis and liver failure before adulthood. It is characterized by elevated serum gamma-glutamyltransferase levels. Defects in ABCB4 are a cause of intrahepatic cholestasis of pregnancy (ICP) [MIM:147480]; also known as obstetric cholestasis. ICP is a multifactorial liver disorder of pregnancy. It presents during the second or, more commonly, the third trimestre of pregnancy with intense pruritus which becomes more severe with advancing gestation and cholestasis. Cholestasis results from abnormal biliary transport from the liver into the small intestine. ICP causes fetal distress, spontaneous premature delivery and intrauterine death. ICP patients have spontaneous and progressive disappearance of cholestasis after delivery. Defects in ABCB4 are a cause of gallbladder disease type 1 (GBD1) [MIM:600803]. It is one of the major digestive diseases. Gallstones composed of cholesterol (cholelithiasis) are the common manifestations in western countries. Most people with gallstones, however, remain asymptomatic through their lifetimes.
配列類似性	Belongs to the ABC transporter superfamily. ABCB family. Multidrug resistance exporter (TC 3.A.1.201) subfamily. Contains 2 ABC transmembrane type-1 domains. Contains 2 ABC transporter domains.
細胞内局在	Cell membrane.

画像



Overlay histogram showing HeLa cells stained with ab24108 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab24108, 1µg/1x10⁶ cells) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-mouse IgG (H+L) ([ab96879](#)) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was mouse IgG2b [PLPV219] ([ab91366](#), 2µg/1x10⁶ cells) used under the same conditions. Acquisition of >5,000 events was performed. This antibody gave a positive signal in HeLa cells fixed with 4% paraformaldehyde (10 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors