

### Anti-ACADVL/VLCAD antibody [EPR15107(B)] ab188872

KO 評価済 リコンビナント RabMAb

★★★★★ **1 Abreviews** 画像数 8

#### 製品の概要

製品名	Anti-ACADVL/VLCAD antibody [EPR15107(B)]
製品の詳細	Rabbit monoclonal [EPR15107(B)] to ACADVL/VLCAD
由来種	Rabbit
アプリケーション	<b>適用あり:</b> Flow Cyt (Intra), IP, WB, IHC-P
種交差性	<b>交差種:</b> Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: HeLa, HepG2, A431 and 293 cell lysates. IHC-P: Human kidney tissue and colon adenocarcinoma tissue. WB: HeLa cells.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"><li>- High batch-to-batch consistency and reproducibility</li><li>- Improved sensitivity and specificity</li><li>- Long-term security of supply</li><li>- Animal-free production</li></ul> <p>For more information <a href="#">see here</a>.</p> <p>Our RabMAb<sup>®</sup> technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to <a href="#">RabMAb<sup>®</sup> patents</a>.</p>

#### 製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 40% Glycerol (glycerin, glycerine), 59% PBS, 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR15107(B)
アイソタイプ	IgG

## アプリケーション


**The Abpromise guarantee** Abpromise保証は、次のテスト済みアプリケーションにおけるab188872の使用に適用されず  
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		1/160. <b>ab172730</b> - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
IP		1/40.
WB		1/1000 - 1/10000. Detects a band of approximately 66 kDa (predicted molecular weight: 70 kDa).
IHC-P		1/250 - 1/500. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

## ターゲット情報

機能	Active toward esters of long-chain and very long chain fatty acids such as palmitoyl-CoA, mysritoyl-CoA and stearoyl-CoA. Can accomodate substrate acyl chain lengths as long as 24 carbons, but shows little activity for substrates of less than 12 carbons.
パスウェイ	Lipid metabolism; mitochondrial fatty acid beta-oxidation.
関連疾患	Defects in ACADVL are the cause of acyl-CoA dehydrogenase very long chain deficiency (ACADVLD) [MIM:201475]. ACADVLD is an autosomal recessive disease which leads to impaired long-chain fatty acid beta-oxidation. It is clinically heterogeneous, with three major phenotypes: a severe childhood form, with early onset, high mortality, and high incidence of cardiomyopathy; a milder childhood form, with later onset, usually with hypoketotic hypoglycemia as the main presenting feature, low mortality, and rare cardiomyopathy; and an adult form, with isolated skeletal muscle involvement, rhabdomyolysis, and myoglobinuria, usually triggered by exercise or fasting.
配列類似性	Belongs to the acyl-CoA dehydrogenase family.
細胞内局在	Mitochondrion inner membrane.

## 画像



Western blot - Anti-ACADVL/VLCAD antibody [EPR15107(B)] (ab188872)

**All lanes :** Anti-ACADVL/VLCAD antibody [EPR15107(B)] (ab188872) at 1/1000 dilution

**Lane 1 :** Wild-type HEK293T cell lysate

**Lane 2 :** ACADVL knockout HEK293T cell lysate

**Lane 3 :** A431 cell lysate

**Lane 4 :** HepG2 cell lysate

Lysates/proteins at 20 µg per lane.

### Secondary


**All lanes :** Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) at 1/10000 dilution

**Predicted band size:** 70 kDa

**Observed band size:** 70 kDa


**Lanes 1-4:** Merged signal (red and green). Green - ab188872 observed at 70 kDa. Red - loading control [ab8245](#) observed at 36 kDa.

ab188872 Anti-ACADVL/VLCAD antibody [EPR15107(B)] was shown to specifically react with ACADVL/VLCAD in wild-type HEK293T cells. Loss of signal was observed when knockout cell line [ab266484](#) (knockout cell lysate [ab257332](#)) was used. Wild-type and ACADVL/VLCAD knockout samples were subjected to SDS-PAGE. ab188872 and Anti-GAPDH antibody [6C5] - Loading Control ([ab8245](#)) were incubated overnight at 4°C at 1 in 1000 dilution and 1 in 20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preadsorbed ([ab216773](#)) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preadsorbed ([ab216776](#)) secondary antibodies at 1 in 20000 dilution for 1 hour at room temperature before imaging.



Flow Cytometry (Intracellular) - Anti-ACADVL/VLCAD antibody [EPR15107(B)] (ab188872)


Intracellular flow cytometric analysis of HeLa cells (paraformaldehyde-fixed, 2%) labeling ACADVL/VLCAD with ab188872 at 1/160 dilution (red) or a rabbit IgG (negative) (green), followed by Goat anti rabbit IgG (FITC) secondary at 1/150 dilution.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ACADVL/VLCAD antibody [EPR15107(B)] (ab188872)

Immunohistochemical analysis of paraffin-embedded Human colon adenocarcinoma tissue labeling ACADVL/VLCAD with ab188872 at 1/500 dilution followed by pre-diluted HRP Polymer for Rabbit IgG secondary antibody and counter-stained with Hematoxylin.


Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ACADVL/VLCAD antibody [EPR15107(B)] (ab188872)

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling ACADVL/VLCAD with ab188872 at 1/500 dilution followed by pre-diluted HRP Polymer for Rabbit IgG secondary antibody and counter-stained with Hematoxylin.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Western blot - Anti-ACADVL/VLCAD antibody [EPR15107(B)] (ab188872)

Anti-ACADVL/VLCAD antibody [EPR15107(B)] (ab188872) at 1/1000 dilution + 293 cell lysate at 20 µg

**Secondary**

Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at 1/1000 dilution


**Predicted band size:** 70 kDa



Immunoprecipitation - Anti-ACADVL/VLCAD antibody [EPR15107(B)] (ab188872)

Western blot analysis of immunoprecipitation pellet from A431 lysate immunoprecipitated using ab188872 at 1/50 dilution.

Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at 1/1000 dilution.



Western blot - Anti-ACADVL/VLCAD antibody [EPR15107(B)] (ab188872)

**All lanes :** Anti-ACADVL/VLCAD antibody [EPR15107(B)] (ab188872) at 1/10000 dilution

**Lane 1 :** HeLa cell lysate

**Lane 2 :** HepG2 cell lysate

**Lane 3 :** A431 cell lysate

Lysates/proteins at 20 µg per lane.

**Secondary**

**All lanes :** Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugate at 1/1000 dilution

**Predicted band size:** 70 kDa

### Why choose a recombinant antibody?



**Research with confidence**  
Consistent and reproducible results



**Long-term and scalable supply**  
Recombinant technology



**Success from the first experiment**  
Confirmed specificity



**Ethical standards compliant**  
Animal-free production

Anti-ACADVL/VLCAD antibody [EPR15107(B)]  
(ab188872)

**Please note:** All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

---

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

### Terms and conditions

---

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors