

Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971] ab109368

KO 評価済 リコンビナント RabMAb

6 References 画像数 11

製品の概要

製品名	Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971]
製品の詳細	Rabbit monoclonal [EPR4971] to Acetyl Coenzyme A Carboxylase
由来種	Rabbit
アプリケーション	適用あり: Flow Cyt (Intra), WB, IHC-P, ICC/IF 適用なし: IP
種交差性	交差種: Human 非交差種: Mouse, Rat
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	HepG2, and SH-SY5Y cell lysates. Human brain tissue and Human skeletal muscle tissue. 293T cells.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Avoid freeze / thaw cycle. Stable for 12 months at -20°C.
バッファー	pH: 7.20 Preservative: 0.01% Sodium azide

	Constituents: 59% PBS, 40% Glycerol, 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR4971
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab109368の使用に適用されず**
 アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。


アプリケーション	Abreviews	特記事項
Flow Cyt (Intra)		1/100 - 1/500. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.
WB		1/5000. Predicted molecular weight: 266 kDa. For unpurified use at 1/1000- 1/10000.
IHC-P		1/250 - 1/500. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
ICC/IF		1/100 - 1/250.

追加情報 Is unsuitable for IP.

ターゲット情報

機能	Catalyzes the rate-limiting reaction in the biogenesis of long-chain fatty acids. Carries out three functions: biotin carboxyl carrier protein, biotin carboxylase and carboxyltransferase.
組織特異性	Expressed in brain, placental, skeletal muscle, renal, pancreatic and adipose tissues; expressed at low level in pulmonary tissue; not detected in the liver.
パスウェイ	Lipid metabolism; malonyl-CoA biosynthesis; malonyl-CoA from acetyl-CoA: step 1/1.
関連疾患	Acetyl-CoA carboxylase 1 deficiency
配列類似性	Contains 1 ATP-grasp domain. Contains 1 biotin carboxylation domain. Contains 1 biotinyl-binding domain. Contains 1 carboxyltransferase domain.
翻訳後修飾	Phosphorylation on Ser-1263 is required for interaction with BRCA1.
細胞内局在	Cytoplasm.

画像



Western blot - Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971] (ab109368)

All lanes : Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971] (ab109368) at 1/1000 dilution

Lane 1 : Wild-type HAP1 whole cell lysate

Lane 2 : ACACA (Acetyl Coenzyme A Carboxylase) knockout HAP1 whole cell lysate

Lane 3 : HeLa whole cell lysate


Lane 4 : A431 whole cell lysate

Lysates/proteins at 20 µg per lane.

Predicted band size: 266 kDa


Lanes 1 - 4: Merged signal (red and green). Green - ab109368 observed at 265 kDa. Red - loading control, **ab130007**, observed at 130 kDa.

ab109368 was shown to specifically react with Acetyl Coenzyme A carboxylase in wild-type HAP1 cells as signal was lost in ACACA (Acetyl Coenzyme A Carboxylase) knockout cells. Wild-type and ACACA (Acetyl Coenzyme A Carboxylase) knockout samples were subjected to SDS-PAGE. Ab109368 and **ab130007** (Mouse anti-Vinculin loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed **ab216773** and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed **ab216776** secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.




Immunocytochemistry/ Immunofluorescence - Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971] (ab109368)

Immunocytochemistry/ Immunofluorescence analysis of 293 (Human embryonic kidney epithelial cell) cells labeling Acetyl Coenzyme A carboxylase with Purified ab109368 at 1:250 dilution (2.1 µg/ml). Cells were fixed in 4% Paraformaldehyde and permeabilized with 0.1% tritonX-100. Cells were counterstained with Ab195889 Anti-alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594) 1:200 (2.5 µg/ml). **ab150077** Goat anti rabbit IgG(Alexa Fluor® 488) was used as the secondary antibody at 1:1000 dilution. DAPI nuclear counterstain. PBS instead of the primary antibody was used as the secondary antibody only control.




Flow Cytometry (Intracellular) - Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971] (ab109368)

Intracellular Flow Cytometry analysis of SH-SY5Y (Human neuroblastoma epithelial cell) cells labeling Acetyl Coenzyme A carboxylase with purified ab109368 at 1/100 dilution (5 µg/ml) (red). Cells were fixed with 4% Paraformaldehyde. A Goat anti rabbit IgG (Alexa Fluor® 488) secondary antibody was used at 1/2000 dilution. Isotype control - 90% methanol. Unlabeled control - Rabbit monoclonal IgG (Black). Cells without incubation with primary antibody and secondary antibody (Blue).




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971] (ab109368)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human kidney tissue sections labeling Acetyl Coenzyme A carboxylase with purified ab109368 at 1:500 dilution (1.05 µg/ml). Heat mediated antigen retrieval was performed using citrate Buffer, pH6.0. Tissue was counterstained with Hematoxylin. **ab97051** Goat Anti-Rabbit IgG H&L (HRP) secondary antibody was used at 1:500 dilution. PBS instead of the primary antibody was used as the negative control.



Flow Cytometry (Intracellular) - Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971] (ab109368)

Overlay histogram showing SH-SY5Y cells stained with unpurified ab109368 (red line). The cells were fixed with 4% paraformaldehyde (10 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab109368, 1/1000 dilution) for 30 min at 22°C. The secondary antibody used was DyLight® 488 goat anti-rabbit IgG (H+L) (**ab96899**) at 1/500 dilution for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) (0.1µg/1x10⁶ cells) used under the same conditions. Unlabelled sample (blue line) was also used as a control. Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter. This antibody gave a positive signal in SH-SY5Y cells fixed with 80% methanol (5 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.



Western blot - Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971] (ab109368)

All lanes : Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971] (ab109368) at 1/1000 dilution (unpurified)

Lane 1 : 293T cell lysate

Lane 2 : HepG2 cell lysate


Lane 3 : SH-SY5Y cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : HRP labelled goat anti-rabbit at 1/2000 dilution

Predicted band size: 266 kDa



Immunocytochemistry/ Immunofluorescence - Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971] (ab109368)

Immunofluorescent staining of 293 cells using unpurified ab109368 at 1/100 dilution



Western blot - Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971] (ab109368)

All lanes : Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971] (ab109368) at 1/5000 dilution

Lane 1 : 293 (Human embryonic kidney epithelial cell) whole cell lysate

Lane 2 : K562 (Human chronic myelogenous leukemia lymphoblast) whole cell lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution

Predicted band size: 266 kDa

Observed band size: 266 kDa


5% NFDm/TBST



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971] (ab109368)

Immunohistochemical analysis of paraffin-embedded skeletal muscle tissue using unpurified ab109368 at 1/250 dilution.

Heat mediated antigen retrieval was performed with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971] (ab109368)

Immunohistochemical analysis of paraffin-embedded brain tissue using unpurified ab109368 at 1/250 dilution.

Heat mediated antigen retrieval was performed with citrate buffer pH 6 before commencing with IHC staining protocol.

Why choose a recombinant antibody?



Anti-Acetyl Coenzyme A Carboxylase antibody [EPR4971] (ab109368)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors