abcam

Product datasheet

Anti-ADH5 antibody [EPR12885(B)] ab177932

יולצעבע RabMAb

★★★★★ 3 Abreviews 3 References 画像数7

製品の概要

製品名 Anti-ADH5 antibody [EPR12885(B)]

製品の詳細 Rabbit monoclonal [EPR12885(B)] to ADH5

由来種 Rabbit

特異性 The mouse and rat recommendation is based on the WB results. We do not guarantee IHC-P for

mouse and rat.

アプリケーション 適用あり: WB, IHC-P

適用なし: Flow Cyt,ICC/IF or IP

種交差性 交差種: Mouse, Rat, Human

免疫原 Synthetic peptide within Human ADH5 aa 1-100. The exact sequence is proprietary.

Database link: P11766

ポジティブ・コントロール WB: Human liver, Mouse liver, Human fetal liver, HepG2, Human fetal kidney, SH-SY5Y, Mouse

brain, Rat liver, and Rat brain lysates. IHC-P: human kidney tissue.

特記事項 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

バッファー Preservative: 0.01% Sodium azide

Constituents: PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA

精製度 Protein A purified

ポリモノ モノクローナル

クローン名

EPR12885(B)

アイソタイプ

lgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab177932の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (2)	1/1000 - 1/5000. Predicted molecular weight: 40 kDa.
IHC-P		1/8000. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol. The mouse and rat recommendation is based on the WB results. We do not guarantee IHC-P for mouse and rat. See IHC antigen retrieval protocols. For unpurified use at 1/50-1/100

追加情報


Is unsuitable for Flow Cyt,ICC/IF or IP.

ターゲット情報

機能	Class-III ADH is remarkably ineffective in oxidizing ethanol, but it readily catalyzes the oxidation of	
	long-chain primary alcohols and the oxidation of S-(hydroxymethyl) glutathione.	
配列類似性	Belongs to the zinc-containing alcohol dehydrogenase family. Class-Ill subfamily.	

細胞内局在 Cytoplasm.

画像




Western blot - Anti-ADH5 antibody [EPR12885(B)] (ab177932)

Anti-ADH5 antibody [EPR12885(B)] (ab177932) at 1/5000 dilution (Purified) + Human liver lysates at 15 μg

Secondary

Goat Anti-Rabbit $\lg G$ (HRP) with minimal cross-reactivity with human $\lg G$ at 1/2000 dilution


Predicted band size: 40 kDa **Observed band size:** 40 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-ADH5 antibody

[EPR12885(B)] (ab177932)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of human kidney tissue sections labeling ADH5 with purified ab177932 at 1/8000 dilution (0.06 µg/ml). Perform heat mediated antigen retrieval using ab93684 (Tris/EDTA buffer, pH 9.0). ImmunoHistoProbe one step HRP Polymer (ready to use) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.



Western blot - Anti-ADH5 antibody [EPR12885(B)] (ab177932)

All lanes : Anti-ADH5 antibody [EPR12885(B)] (ab177932) at 1/5000 dilution (Purified)

Lane 1 : Mouse liver lysates
Lane 2 : Mouse brain lysates
Lane 3 : Rat liver lysates
Lane 4 : Rat brain lysates


Lysates/proteins at 15 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit lgG H&L (HRP) (ab97051) at 1/20000 dilution

Predicted band size: 40 kDa **Observed band size:** 40 kDa

ADH5 is highly expressed in liver, kidney and low expressed in brain as is described in PMID: 26412304



Western blot - Anti-ADH5 antibody [EPR12885(B)] (ab177932)

All lanes : Anti-ADH5 antibody [EPR12885(B)] (ab177932) at 1/1000 dilution ((unpurified))

Lane 1: Human fetal liver lysates


Lane 2: HepG2 lysates

Lane 3: Human fetal kidney lysates

Lane 4: SH-SY5Y lysates

Lysates/proteins at 10 µg per lane.


Predicted band size: 40 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-ADH5 antibody
[EPR12885(B)] (ab177932)

Immunohistochemical analysis of paraffin-embedded Human kidney tissue labeling ADH5 with unpurified ab177932 at a 1/50 dilution.


Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-ADH5 antibody
[EPR12885(B)] (ab177932)

Immunohistochemical analysis of paraffin-embedded Human brain tissue labeling ADH5 with unpurified ab177932 at a 1/50 dilution.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors