abcam

Product datasheet

Alexa Fluor® 488 Anti-NeuN antibody [EPR12763] - Neuronal Marker ab190195

יובעבלי RabMAb

★★★★★ 6 Abreviews 33 References

画像数 5

製品名 Alexa Fluor® 488 Anti-NeuN antibody [EPR12763] - Neuronal Marker

製品の詳細 Alexa Fluor® 488 Rabbit monoclonal [EPR12763] to NeuN - Neuronal Marker

由来種 Rabbit

Alexa Fluor® 488. Ex: 495nm, Em: 519nm 標識

アプリケーション 適用あり: ICC/IF, IHC-Fr, Flow Cyt (Intra)

種交差性 交差種: Rat, Human

交差が予測される動物種: Mouse, Sheep, Goat, Cat, Dog, Zebrafish, Common marmoset

Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ICC/IF: NGF-differentiated PC12 cells and U-87 MG cells. IHC-Fr: Rat Brain (Normal). Flow Cyt

(intra): U-87 MG cells.

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor[®] dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact

製品の概要

免疫原

ポジティブ・コントロール

特記事項

Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle. Stable for 12 months at -20°C. Store In the Dark.

バッファー pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA

精製度 Protein A purified

ポリモノ モノクローナル **クローン名** EPR12763

アイソタイプ IgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab190195の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF	★★★★ (1)	1/50 - 1/250.
IHC-Fr	★★★★☆ (1)	1/50. Before commencing with immunostaining protocol, perform heat mediated antigen retrieval using sodium citrate buffer, pH6.
Flow Cyt (Intra)		1/500. ab199091 - Rabbit monoclonal IgG (Alexa Fluor® 488), is suitable for use as an isotype control with this antibody.


ターゲット情報

機能 RNA-binding protein that regulates alternative splicing events.

配列類似性 Contains 1 RRM (RNA recognition motif) domain.

細胞内局在 Nucleus. Cytoplasm.

画像




Immunohistochemistry (Frozen sections) - Alexa Fluor® 488 Anti-NeuN antibody [EPR12763] -Neuronal Marker (ab190195)

IHC image of ab190195 staining in acetone fixed frozen tissue section of normal rat brain.

Non-specific protein-protein interactions were blocked using TBS containing 0.025% (v/v) Triton X-100, 0.3M (w/v) glycine and 3% (w/v) BSA for 1h at room temperature. The section was then incubated with ab190195 (1/50) in TBS containing 0.025% (v/v) Triton X-100 and 3% (w/v) BSA overnight at +4°C. The section was then counterstained and mounted with SlowFade® Gold Antifade Mountant with DAPI.


The DAPI only control (no antibody) inset shows no autofluorescence, demonstrating that any Alexa Fluor[®] 488 signal is dervied directly from bound ab190195. The separate images of ab190195 and DAPI alone, combined with the merged version of both signals, shows predominant co-localisation of the Alexa Fluor[®] 488 signal in the nuclei of the hippocampal granular layer.

For other IHC staining systems (automated and non-automated), customers should optimize variable parameters such as antigen retrieval conditions, antibody concentrations and incubation times.



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 488 Anti-NeuN antibody [EPR12763] -Neuronal Marker (ab190195)

ab190195 staining NeuN in U87-MG cells. The cells were fixed with 100% methanol (5min) and then blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab190195 at 1/50 dilution (shown in green) and ab7291 (Mouse monoclonal [DM1A] to alpha Tubulin) at 1 μ g/ml overnight at +4°C, followed by a further incubation at room temperature for 1h with an Alexa Fluor 594 Goat anti-Mouse secondary (ab150120) at 2 μ g/ml (shown in red). Nuclear DNA was labelled in blue with DAPI.



Flow Cytometry (Intracellular) - Alexa Fluor® 488 Anti-NeuN antibody [EPR12763] - Neuronal Marker (ab190195)

ab 190195 ab 7231


Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 488 Anti-NeuN antibody [EPR12763] -Neuronal Marker (ab190195)

Overlay histogram showing U-87MG cells stained with ab190195 (red line). The cells were fixed with 80% methanol (5 min) and then permeabilized with 0.1% PBS-Tween for 20 min. The cells were then incubated in 1x PBS / 10% normal goat serum / 0.3M glycine to block non-specific protein-protein interactions followed by the antibody (ab190195, 1/500 dilution) for 30 min at 22°C. Isotype control antibody (black line) was rabbit IgG (monoclonal) Alexa Fluor® 488 used at the same concentration and conditions as the primary antibody. Unlabelled sample (blue line) was also used as a control.

Acquisition of >5,000 events were collected using a 20mW Argon ion laser (488nm) and 525/30 bandpass filter.

This antibody gave a positive signal in U-87MG fixed with 4% formaldehyde (10 min)/permeabilized with 0.1% PBS-Tween for 20 min used under the same conditions.

ab190195 staining NeuN in NGF-differentiated PC12 cells (7 days). The cells were fixed with 100% methanol (5min) and then blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated with ab190195 at 1/50 dilution (shown in green) and ab7291 (Mouse monoclonal [DM1A] to alpha Tubulin) at 1 μ g/ml overnight at +4°C, followed by a further incubation at room temperature for 1h with an Alexa Fluor[®] 594 Goat anti-Mouse secondary (ab150120) at 2 μ g/ml (shown in red). Nuclear DNA was labelled in blue with DAPI.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors