abcam

Product datasheet

Alexa Fluor® 568 Anti-Cytokeratin 19 antibody [EP1580Y] ab203445

יובעבלי RabMAb

2 References 画像数 2

製品の概要

特記事項

製品名 Alexa Fluor® 568 Anti-Cytokeratin 19 antibody [EP1580Y]

製品の詳細 Alexa Fluor® 568 Rabbit monoclonal [EP1580Y] to Cytokeratin 19

由来種 Rabbit

Alexa Fluor® 568. Ex: 578nm, Em: 603nm 標識

アプリケーション 適用あり: ICC/IF 種交差性 交差種: Human

交差が予測される動物種: Mouse

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール ICC/IF: HeLa cells

> Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb® patents.

Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or

outlicensing@thermofisher.com.

製品の特性

製品の状態 Liquid


保存方法

Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

解離定数(K_D値)

 $K_D = 3.70 \times 10^{-10} M$



Learn more about K_D

バッファー pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA

精製度 Protein A purified

ポリモノクローナル **ウローン名** EP1580Y

アイソタイプ IgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab203445の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		1/200. This product gave a positive signal in HeLa cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min)

ターゲット情報

機能 Involved in the organization of myofibers. Together with KRT8, helps to link the contractile

apparatus to dystrophin at the costameres of striated muscle.

組織特異性 Expressed in a defined zone of basal keratinocytes in the deep outer root sheath of hair follicles.

Also observed in sweat gland and mammary gland ductal and secretory cells, bile ducts, gastrointestinal tract, bladder urothelium, oral epithelia, esophagus, ectocervical epithelium (at protein level). Expressed in epidermal basal cells, in nipple epidermis and a defined region of the hair follicle. Also seen in a subset of vascular wall cells in both the veins and artery of human umbilical cord, and in umbilical cord vascular smooth muscle. Observed in muscle fibers accumulating in the costameres of myoplasm at the sarcolemma in structures that contain


dystrophin and spectrin.

配列類似性 Belongs to the intermediate filament family.

発生段階 Present in hair follicles at all stages of development.

ドメイン This keratin differs from all other IF proteins in lacking the C-terminal tail domain.

画像



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 568 Anti-Cytokeratin 19 antibody [EP1580Y] (ab203445)

ab203445 staining Cytokeratin 19 in HeLa cells. The cells were fixed with 100% methanol (5min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab203445 at 1/200 dilution (pseudocolored in green) and ab195884, Rat monoclonal to Tubulin (Alexa Fluor[®] 647), at 1/250 dilution (shown in red). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

This product also gave a positive signal under the same testing conditions in HeLa cells fixed with 4% formaldehyde (10 min).



[EP1580Y] (ab203445)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- · Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors