abcam

Product datasheet

Alexa Fluor® 647 Anti-ATPB antibody [3D5] - Mitochondrial Marker ab197649

1 References 画像数 1

製品の概要

製品名 Alexa Fluor® 647 Anti-ATPB antibody [3D5] - Mitochondrial Marker

製品の詳細 Alexa Fluor® 647 Mouse monoclonal [3D5] to ATPB - Mitochondrial Marker

由来種 Mouse

標識 Alexa Fluor® 647. Ex: 652nm, Em: 668nm

 アプリケーション
 適用あり: ICC/IF

 種交差性
 交差種: Human

交差が予測される動物種: Mouse, Rat, Cow, Caenorhabditis elegans, Monkey

A

The details of the immunogen for this antibody are not available.

ICC/IF: HepG2 cells.

Alexa Fluor[®] is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor[®] dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor[®] dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor[®] dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or **outlicensing@thermofisher.com**.

The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.

If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As

免疫原

ポジティブ・コントロール

特記事項

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C or -

80°C. Avoid freeze / thaw cycle. Store In the Dark.

バッファー pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: PBS, 30% Glycerol (glycerin, glycerine), 1% BSA

精製度 IgG fraction

特記事項(精製) Near homogeneity as judged by SDS-PAGE. The antibody was produced in vitro using

hybridomas grown in serum-free medium, and then purified by biochemical fractionation.

ポリ/モノ モノクローナル

クローン名3D5アイソタイプIgG1軽鎖の種類kappa

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab197649の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		1/50. This product gave a positive signal in HepG2 cells fixed with 100% methanol (5 min).

ターゲット情報

機能 Mitochondrial membrane ATP synthase (F(1)F(0) ATP synthase or Complex V) produces ATP


from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F(1) - containing the extramembraneous catalytic core, and F(0) - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Subunits alpha and beta form the catalytic core in F(1). Rotation of the central stalk against the surrounding alpha(3)beta(3) subunits leads to

hydrolysis of ATP in three separate catalytic sites on the beta subunits.

配列類似性 Belongs to the ATPase alpha/beta chains family.

細胞内局在 Mitochondrion. Mitochondrion inner membrane. Peripheral membrane protein.

画像



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 647 Anti-ATPB antibody [3D5] -Mitochondrial Marker (ab197649) ab197649 staining ATPB in HepG2 cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab197649 at a 1/50 dilution (shown in red) and ab195887, Mouse monoclonal to alpha Tubulin (Alexa Fluor[®] 488), at a 1/250 dilution (shown in green). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- · Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

· Guarantee only valid for products bought direct from Abcam or one of our authorized distributors