# abcam

## Product datasheet

## Alexa Fluor® 647 Anti-Hexokinase 1 antibody [EPR10134(B)] -Mitochondrial Outer Membrane Marker ab197864

יובעדער RabMAb

★★★★★ 1 Abreviews 1 References

画像数 2

#### 製品の概要

製品名 Alexa Fluor® 647 Anti-Hexokinase 1 antibody [EPR10134(B)] - Mitochondrial Outer Membrane

Marker

製品の詳細 Alexa Fluor® 647 Rabbit monoclonal [EPR10134(B)] to Hexokinase 1 - Mitochondrial Outer

Membrane Marker

由来種 Rabbit

標識 Alexa Fluor® 647. Ex: 652nm. Em: 668nm

アプリケーション 適用あり: ICC/IF 種交差性 交差種: Human

交差が予測される動物種: Mouse, Rat 🔷

免疫原 Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール ICC/IF: HCT116 cells.

特記事項 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dye, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor® dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research.

1

For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or <a href="mailto:outlicensing@thermofisher.com">outlicensing@thermofisher.com</a>.

#### 製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

**バッファー** pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, PBS

精製度 Protein A purified

**ポリ/モノ** モノクローナル **クローン名** EPR10134(B)

アイソタイプ lgG

#### アプリケーション

**The Abpromise guarantee** <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab197864の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		1/50. This product gave a positive signal in HCT116 cells fixed with 100% methanol (5 min).

#### ターゲット情報

組織特異性 lsoform 2 is erythrocyte specific. lsoform 3 and isoform 4 are testis-specific.

אליכל Carbohydrate metabolism; hexose metabolism.

関連疾患 Hexokinase deficiency

Neuropathy, hereditary motor and sensory, Russe type

**配列類似性** Belongs to the hexokinase family.

Contains 2 hexokinase domains.

The N- and C-terminal halves of this hexokinase show extensive sequence similarity to each


other. The catalytic activity is associated with the C-terminus while regulatory function is

associated with the N-terminus. Each domain can bind a single glucose and Gluc-6-P molecule.

細胞内局在 Mitochondrion outer membrane. Its hydrophobic N-terminal sequence may be involved in


membrane binding.

#### 画像



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 647 Anti-Hexokinase 1 antibody [EPR10134(B)] - Mitochondrial Outer Membrane Marker (ab197864) ab197864 staining Hexokinase 1 in HCT116 cells. The cells were fixed with 100% methanol (5 min), permeabilized with 0.1% Triton X-100 for 5 minutes and then blocked with 1% BSA/10% normal goat serum/0.3M glycine in 0.1%PBS-Tween for 1h. The cells were then incubated overnight at +4°C with ab at a 1/50 dilution (shown in red) and <a href="mailto:ab195887">ab195887</a>, Mouse monoclonal to alpha Tubulin (Alexa Fluor<sup>®</sup> 488), at a 1/250 dilution (shown in green). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).



Marker (ab197864)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

### Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <a href="https://www.abcam.co.jp/abpromise">https://www.abcam.co.jp/abpromise</a> or contact our technical team.

## Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors