abcam

Product datasheet

Alexa Fluor® 647 Anti-PKM antibody [EPR10138(B)] ab214257

ועלשעבע RabMAb

★★★★ 1 Abreviews 画像数 2

製品の概要

製品名 Alexa Fluor® 647 Anti-PKM antibody [EPR10138(B)]

製品の詳細 Alexa Fluor® 647 Rabbit monoclonal [EPR10138(B)] to PKM

由来種 Rabbit

標識 Alexa Fluor® 647. Ex: 652nm. Em: 668nm

アプリケーション 適用あり: ICC/IF 種交差性 交差種: Human

Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.

交差が予測される動物種: Mouse, Rat, Pig

ICC/IF: HeLa cells

This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility
- Improved sensitivity and specificity
- Long-term security of supply
- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb**® **patents**.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc, a Thermo Fisher Scientific Company. The Alexa Fluor® dye included in this product is provided under an intellectual property license from Life Technologies Corporation. As this product contains the Alexa Fluor® dve, the purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). As this product contains the Alexa Fluor[®] dye the sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: in manufacturing; (ii) to provide a service, information, or data in return for payment (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are sold for use in research. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

免疫原

ポジティブ・コントロール

特記事項

製品の特性

製品の状態 Liquid

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C.

Avoid freeze / thaw cycle. Store In the Dark.

バッファー pH: 7.40

Preservative: 0.02% Sodium azide

Constituents: 30% Glycerol (glycerin, glycerine), 1% BSA, PBS

精製度 Protein A purified

ポリ/モノ モノクローナル **クローン名** EPR10138(B)

アイソタイプ IgG

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab214257の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		1/100. This product gave a positive signal in HeLa cells fixed with 4% formaldehyde (10 min) and 100% methanol (5 min)

ターゲット情報

機能
Glycolytic enzyme that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate
(PEP) to ADP, generating ATP. Stimulates POU5F1-mediated transcriptional activation. Plays a
general role in caspase independent cell death of tumor cells. The ratio between the highly
active tetrameric form and nearly inactive dimeric form determines whether glucose carbons are
channeled to biosynthetic processes or used for glycolytic ATP production. The transition between

survival.

組織特異性 Specifically expressed in proliferating cells, such as embryonic stem cells, embryonic carcinoma

cells, as well as cancer cells.

パスウェイ Carbohydrate degradation; glycolysis; pyruvate from D-glyceraldehyde 3-phosphate: step 5/5.

配列類似性 Belongs to the pyruvate kinase family.

翻訳後修飾 ISGylated.

Under hypoxia, hydroxylated by EGLN3.


Acetylation at Lys-305 is stimulated by high glucose concentration, it decreases enzyme activity and promotes its lysosomal-dependent degradation via chaperone-mediated autophagy. FGFR1-dependent tyrosine phosphorylation is reduced by interaction with TRIM35.

the 2 forms contributes to the control of glycolysis and is important for tumor cell proliferation and

細胞内局在 Cytoplasm. Nucleus. Translocates to the nucleus in response to different apoptotic stimuli.

Nuclear translocation is sufficient to induce cell death that is caspase independent, isoform-


画像



Immunocytochemistry/ Immunofluorescence - Alexa Fluor® 647 Anti-PKM antibody [EPR10138(B)] (ab214257) Ab214257 staining PKM in HeLa cells. The cells were fixed with 4% formaldehyde (10 min), and then incubated in 1%BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween for 1h to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated overnight at +4°C with ab214257 at 1/50 dilution (shown in red) and ab195887, Mouse monoclonal [DM1A] to alpha Tubulin - Microtubule Marker (Alexa Fluor® 488), at 1/250 dilution (shown in green). Nuclear DNA was labelled with DAPI (shown in blue).

Image was taken with a confocal microscope (Leica-Microsystems, TCS SP8).

This product also gave a positive signal under the same testing conditions in HeLa cells fixed with 100% methanol (5min).



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

• Guarantee only valid for products bought direct from Abcam or one of our authorized distributors