

Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] ab179484

リコンビナント **RabMAb**

★★★★★ **4 Abreviews** **34 References** 画像数 **14**

製品の概要

製品名	Anti-alpha Tubulin (acetyl K40) antibody [EPR16772]
製品の詳細	Rabbit monoclonal [EPR16772] to alpha Tubulin (acetyl K40)
由来種	Rabbit
アプリケーション	適用あり: WB, ICC/IF, IP, IHC-P, Flow Cyt (Intra)
種交差性	交差種: Mouse, Rat, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: HeLa, C6 and NIH/3T3 whole cell lysates (treated with 500 ng/ml Trichostatin A for 4 hours); Mouse brain, kidney and spleen lysates; Rat brain and heart lysates; Human fetal heart and fetal kidney lysates. IHC-P: Human and Mouse cerebral cortex tissue; rat cerebellum tissue. IF: HeLa cells treated with 50 ug/ml Trichostatin A for 4 hours. Flow: HeLa cells treated with 500ng/ml Trichostatin A for 4 hours. IP: HeLa treated with 500 ng/ml Trichostatin A for 4 hours.
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none"> - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	Preservative: 0.01% Sodium azide Constituents: 40% Glycerol (glycerin, glycerine), 0.05% BSA, 59% PBS
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR16772

アプリケーション


The Abpromise guarantee Abpromise保証は、次のテスト済みアプリケーションにおけるab179484の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご確認ください。

アプリケーション	Abreviews	特記事項
WB		1/2000. Detects a band of approximately 52 kDa (predicted molecular weight: 50 kDa).
ICC/IF	★★★★★ (1)	1/500.
IP		1/70.
IHC-P	★★★★★ (3)	1/1000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.
Flow Cyt (Intra)		1/240. ab172730 - Rabbit monoclonal IgG, is suitable for use as an isotype control with this antibody.

ターゲット情報


機能	Tubulin is the major constituent of microtubules. It binds two moles of GTP, one at an exchangeable site on the beta chain and one at a non-exchangeable site on the alpha chain.
配列類似性	Belongs to the tubulin family.
翻訳後修飾	Some glutamate residues at the C-terminus are polyglutamylated. This modification occurs exclusively on glutamate residues and results in polyglutamate chains on the gamma-carboxyl group. Also monoglycylated but not polyglycylated due to the absence of functional TTL10 in human. Monoglycylation is mainly limited to tubulin incorporated into axonemes (cilia and flagella) whereas glutamylation is prevalent in neuronal cells, centrioles, axonemes, and the mitotic spindle. Both modifications can coexist on the same protein on adjacent residues, and lowering glycylation levels increases polyglutamylated, and reciprocally. The precise function of such modifications is still unclear but they regulate the assembly and dynamics of axonemal microtubules. Acetylation of alpha chains at Lys-40 stabilizes microtubules and affects affinity and processivity of microtubule motors. This modification has a role in multiple cellular functions, ranging from cell motility, cell cycle progression or cell differentiation to intracellular trafficking and signaling.
細胞内局在	Cytoplasm > cytoskeleton.

画像



Immunocytochemistry/ Immunofluorescence - Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] (ab179484)

ab179484 stained in Hela cells. Untreated and Trichostatin A treated (50ug/ml, 4 hours) cells were fixed with 4% paraformaldehyde (10min) at room temperature and incubated with PBS containing 10% goat serum, 0.3 M glycine, 1% BSA and 0.1% triton for 1h at room temperature to permeabilise the cells and block non-specific protein-protein interactions. The cells were then incubated with the antibody ab179484 at 1/500 dilution overnight at +4°C. The secondary antibody was **ab150177** used at 1 ug/ml for 1hour at room temperature (colored green). DAPI was used to stain the cell nuclei (colored blue) at a concentration of 1.43µM for 1hour at room temperature.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] (ab179484)

Immunohistochemical analysis of paraffin-embedded Rat cerebellum tissue labeling alpha Tubulin (acetyl K40) with ab179484 at 1/1000 dilution, followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Cytoplasmic staining is observed on Purkinje cells of cerebellum. Counter stained with Hematoxylin.

Negative control: Using PBS instead of primary ab, secondary ab is prediluted HRP Polymer for Rabbit/Mouse IgG.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



All lanes : Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] (ab179484) at 1/2000 dilution

Lane 1 : Mouse brain lysate

Lane 2 : Mouse kidney lysate

Lane 3 : Mouse spleen lysate

Lane 4 : Rat brain lysate

Lane 5 : Rat heart lysate

Lane 6 : Human fetal heart lysate

Lane 7 : Human fetal kidney lysate

Lysates/proteins at 10 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 50 kDa

Observed band size: 52 kDa

Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] (ab179484) at 1/20000 dilution

Lane 1 : HeLa (Human epithelial cells from cervix adenocarcinoma) whole cell lysate treated with 500 ng/ml Trichostatin A for 4 hours

Lane 2 : Untreated HeLa whole cell lysate

Lysates/proteins at 10 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 50 kDa


Observed band size: 52 kDa

Blocking/Dilution buffer: 5% NFDM/TBST.




Immunocytochemistry/ Immunofluorescence - Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] (ab179484)

Ab179484 staining alpha Tubulin in HFF-1 (Human skin fibroblast) cell line by ICC/IF (Immunocytochemistry/Immunofluorescence). The cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% TritonX-100. Samples were incubated with primary antibody at 1:20000 dilution. An AlexaFluor®488 Goat anti-Rabbit (**ab150077**) was used as a secondary antibody at 1:1000 dilution. An Anti-Alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594), **ab195889** was used as a counterstain at 1:200 dilution. DAPI was used as a nuclear counterstain. Confocal image showing cilia (arrows) staining in HFF-1 cells treated with starvation for 48 hours.




Immunocytochemistry/ Immunofluorescence - Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] (ab179484)

Ab179484 staining alpha Tubulin in NIH/3T3 (mouse embryonic fibroblast) cell line by ICC/IF (Immunocytochemistry/Immunofluorescence). The cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% TritonX-100. Samples were incubated with primary antibody at 1:20000 dilution. An AlexaFluor®488 Goat anti-Rabbit (**ab150077**) was used as a secondary antibody at 1:1000 dilution. An Anti-Alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor® 594), **ab195889** was used as a counterstain at 1:200 dilution. DAPI was used as a nuclear counterstain. Confocal image showing cilia (arrows) staining in NIH/3T3 cells treated with starvation for 48 hours.



Immunocytochemistry/ Immunofluorescence - Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] (ab179484)

Ab179484 staining alpha Tubulin in NIH/3T3 (mouse embryonic fibroblast) cell line by ICC/IF (Immunocytochemistry/Immunofluorescence). The cells were fixed with 4% paraformaldehyde and permeabilized with 0.1% TritonX-100. Samples were incubated with primary antibody at 1:20000 dilution. An AlexaFluor[®]488 Goat anti-Rabbit (**ab150077**) was used as a secondary antibody at 1:1000 dilution. An Anti-Alpha Tubulin antibody [DM1A] - Microtubule Marker (Alexa Fluor[®] 594), **ab195889** was used as a counterstain at 1:200 dilution. DAPI was used as a nuclear counterstain. Confocal image showing midbody (arrows) staining in NIH/3T3 cells treated with starvation for 48 hours.



Western blot - Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] (ab179484)

All lanes : Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] (ab179484) at 1/20000 dilution

Lane 1 : C6 (Rat glial tumor cells) whole cell lysate treated with 500 ng/ml Trichostatin A for 4 hours

Lane 2 : Untreated C6 whole cell lysate

Lysates/proteins at 10 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 50 kDa

Observed band size: 52 kDa

Blocking/Dilution buffer: 5% NFDm/TBST.



All lanes : Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] (ab179484) at 1/20000 dilution

Lane 1 : NIH/3T3 (Mouse embryo fibroblast cells) whole cell lysate treated with 500 ng/ml Trichostatin A for 4 hours

Lane 2 : Untreated NIH/3T3 whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


Lane 1 : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Lane 2 : Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/1000 dilution

Predicted band size: 50 kDa

Additional bands at: 52 kDa. We are unsure as to the identity of these extra bands.

Blocking/dilution buffer: 5% NFDM/TBST.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] (ab179484)

Immunohistochemical analysis of paraffin-embedded Human cerebral cortex tissue labeling alpha Tubulin (acetyl K40) with ab179484 at 1/1000 dilution, followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Cytoplasmic staining is observed on neuron cells of Human brain tissue. Counter stained with Hematoxylin.

Negative control: Using PBS instead of primary ab, secondary ab is prediluted HRP Polymer for Rabbit/Mouse IgG.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] (ab179484)

Immunohistochemical analysis of paraffin-embedded Mouse cerebral cortex tissue labeling alpha Tubulin (acetyl K40) with ab179484 at 1/1000 dilution, followed by prediluted HRP Polymer for Rabbit/Mouse IgG. Cytoplasmic staining is observed on neuron cells of Mouse cerebral cortex tissue. Counter stained with Hematoxylin.


Negative control: Using PBS instead of primary ab, secondary ab is prediluted HRP Polymer for Rabbit/Mouse IgG.

Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Flow Cytometry (Intracellular) - Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] (ab179484)

Intracellular flow cytometric analysis of 2% paraformaldehyde-fixed HeLa (Human epithelial cells from cervix adenocarcinoma) cells treated with 500 ng/ml Trichostatin A for 4 hours labeling alpha Tubulin (acetyl K40) with ab179484 at 1/240 dilution (red line). Goat anti rabbit IgG (FITC) at 1/150 dilution was used as the secondary antibody. ab179484 preincubated with 1 mg/ml acetyl Alpha tubulin (acetyl K40) peptide (green) or non-acetyl Alpha tubulin (acetyl K40) peptide (orange). The isotype control was Rabbit monoclonal IgG (black) and the unlabelled control was cells without incubation with primary antibody and secondary antibody (blue).



Immunoprecipitation - Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] (ab179484)

Alpha Tubulin was immunoprecipitated from 1 mg of HeLa cells (Human epithelial cells from cervix adenocarcinoma) treated with 500 ng/ml Trichostatin A for 4 hours with ab179484 at 1/70 dilution. Western blot was performed from 10 µg of the immunoprecipitate using ab179484 at 1/1000 dilution. Anti-Rabbit IgG (HRP), specific to the non-reduced form of IgG, was used as secondary antibody at 1/1500 dilution. Left lane: Hela whole cell extract. Right lane: PBS instead of Hela whole cell extract.

Blocking and dilution buffer and concentration: 5% NFDM/TBST.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-alpha Tubulin (acetyl K40) antibody [EPR16772] (ab179484)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors