abcam

Product datasheet

Anti-AMH antibody [EPR23380-171] ab272221

ועלשעבע RabMAb

画像数5

製品の概要

製品名 Anti-AMH antibody [EPR23380-171]

製品の詳細 Rabbit monoclonal [EPR23380-171] to AMH

由来種 Rabbit

アプリケーション 適用あり: IHC-Fr, IHC-P

適用なし: IP or WB

種交差性 交差種: Mouse, Rat

非交差種: Human

免疫原 Recombinant fragment. This information is proprietary to Abcam and/or its suppliers.

ポジティブ・コントロール IHC-P: Mouse ovary tissue; Rat ovary tissue. IHC-Fr: Mouse ovary tissue; Rat ovary tissue.

特記事項 This product is a recombinant monoclonal antibody, which offers several advantages including:

- High batch-to-batch consistency and reproducibility

- Improved sensitivity and specificity

- Long-term security of supply

- Animal-free production

For more information see here.

Our RabMAb® technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to **RabMAb patents**.

製品の特性

製品の状態

保存方法 Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long

term. Avoid freeze / thaw cycle.

バッファー pH: 7.2

Preservative: 0.01% Sodium azide

Constituents: PBS, 40% Glycerol, 0.05% BSA

精製度 Protein A purified

ポリモノ モノクローナル

クローン名 EPR23380-171

アプリケーション

The Abpromise guarantee <u>Abpromise保証は、</u>次のテスト済みアプリケーションにおけるab272221の使用に適用されます アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-Fr		1/100. Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).
IHC-P		1/2000. Perform heat mediated antigen retrieval with Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

追加情報

Is unsuitable for IP or WB.

ターゲット情報


関連性

Anti mullerian hormone (AMH) is a member of the TGF beta superfamily. It is secreted as a homodimeric 140kD disulphide linked precursor that is cleaved to release the mature 30kD homodimer. Originally classified as a foetal testicular hormone that inhibits Mullerian duct development, AMH is expressed post natally by immature Sertoli cells, and to a lesser degree by granulosa cells. AMH plays a role in testicular differentiation and in the regulation of ovarian follicle growth.

細胞内局在

Secreted

画像




Immunohistochemistry (Frozen sections) - Anti-AMH antibody [EPR23380-171] (ab272221)

4% PFA-fixed, 0.2% Triton X-100 permeabilised rat ovary tissue labeling AMH with ab272221 at 1/100 dilution followed by <u>ab150077</u> AlexaFluor[®] 488 Goat anti-Rabbit secondary at 1/1000 dilution (Green). Cytoplasmic staining on granulosa cells of follicle. The nuclear counter stain is DAPI (Blue).

Secondary antibody only control: Secondary antibody is **ab150077**AlexaFluor[®]488 Goat anti-Rabbit secondary at 1/1000 dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-AMH antibody
[EPR23380-171] (ab272221)

Immunohistochemical analysis of paraffin-embedded Mouse ovary tissue labeling AMH with ab272221 at 1/2000 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on granulosa cells of mouse ovary. The section was incubated with ab272221 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.




Immunohistochemistry (Frozen sections) - Anti-AMH antibody [EPR23380-171] (ab272221)

4% PFA-fixed, 0.2% Triton X-100 permeabilised mouse ovary tissue labeling AMH with ab272221 at 1/100 dilution followed by <u>ab150077</u> AlexaFluor[®]488 Goat anti-Rabbit secondary at 1/1000 dilution (Green). Cytoplasmic staining on granulosa cells of follicle. The nuclear counter stain is DAPI (Blue).

Secondary antibody only control: Secondary antibody is <u>ab150077</u>
AlexaFluor[®]488 Goat anti-Rabbit secondary at 1/1000 dilution.

Heat mediated antigen retrieval using sodium citrate buffer (10mM citrate pH 6.0 + 0.05% Tween-20).




Immunohistochemistry (Formalin/PFA-fixed paraffinembedded sections) - Anti-AMH antibody
[EPR23380-171] (ab272221)

Immunohistochemical analysis of paraffin-embedded Rat ovary tissue labeling AMH with ab272221 at 1/2000 dilution followed by a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101). Positive staining on granulosa cells of rat ovary. The section was incubated with ab272221 for 30 mins at room temperature. The immunostaining was performed on a Leica Biosystems BOND® RX instrument Counterstained with Hematoxylin.

Secondary antibody only control: Secondary antibody is a ready to use Rabbit specific IHC polymer detection kit HRP/DAB (ab209101).

Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0, epitope retrieval solution2) for 20 mins.



Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- · We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit https://www.abcam.co.jp/abpromise or contact our technical team.

Terms and conditions

Guarantee only valid for products bought direct from Abcam or one of our authorized distributors