

Anti-AMPK gamma 1 antibody [Y307] ab32382

KO 評価済 リコンビナント RabMAb

★★★★★ [1 Abreviews](#) [4 References](#) [画像数 5](#)

製品の概要

製品名	Anti-AMPK gamma 1 antibody [Y307]
製品の詳細	Rabbit monoclonal [Y307] to AMPK gamma 1
由来種	Rabbit
アプリケーション	適用あり: WB, IHC-P
種交差性	交差種: Human
免疫原	Synthetic peptide within Human AMPK gamma 1 aa 1-100. The exact sequence is proprietary.
エピトープ	ab32382 reacts with an epitope located in the N terminal region of AMPK gamma 1.
ポジティブ・コントロール	Jurkat cell lysate and HeLa cell lysate, human skeletal muscles
特記事項	<p>This product is a recombinant monoclonal antibody, which offers several advantages including:</p> <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production <p>For more information see here.</p> <p>Our RabMAb[®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents.</p> <p>Mouse, Rat: We have preliminary internal testing data to indicate this antibody may not react with these species. Please contact us for more information.</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 49% PBS, 50% Glycerol (glycerin, glycerine), 0.05% BSA
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	Y307

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab32382の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★★ (1)	1/1000 - 1/2000. Predicted molecular weight: 38 kDa.
IHC-P		Use at an assay dependent concentration. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.

ターゲット情報

機能	AMP/ATP-binding subunit of AMP-activated protein kinase (AMPK), an energy sensor protein kinase that plays a key role in regulating cellular energy metabolism. In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation. AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators. Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton; probably by indirectly activating myosin. Gamma non-catalytic subunit mediates binding to AMP, ADP and ATP, leading to activate or inhibit AMPK: AMP-binding results in allosteric activation of alpha catalytic subunit (PRKAA1 or PRKAA2) both by inducing phosphorylation and preventing dephosphorylation of catalytic subunits. ADP also stimulates phosphorylation, without stimulating already phosphorylated catalytic subunit. ATP promotes dephosphorylation of catalytic subunit, rendering the AMPK enzyme inactive.
配列類似性	Belongs to the 5'-AMP-activated protein kinase gamma subunit family. Contains 4 CBS domains.
ドメイン	The AMPK pseudosubstrate motif resembles the sequence around sites phosphorylated on target proteins of AMPK, except the presence of a non-phosphorylatable residue in place of Ser. In the absence of AMP this pseudosubstrate sequence may bind to the active site groove on the alpha subunit (PRKAA1 or PRKAA2), preventing phosphorylation by the upstream activating kinase STK11/LKB1. The CBS domains mediate binding to AMP, ADP and ATP. 2 sites bind either AMP or ATP, whereas a third site contains a tightly bound AMP that does not exchange. Under physiological conditions AMPK mainly exists in its inactive form in complex with ATP, which is much more abundant than AMP.
翻訳後修飾	Phosphorylated by ULK1 and ULK2; leading to negatively regulate AMPK activity and suggesting the existence of a regulatory feedback loop between ULK1, ULK2 and AMPK.

画像



Western blot - Anti-AMPK gamma 1 antibody [Y307] (ab32382)

All lanes : Anti-AMPK gamma 1 antibody [Y307] (ab32382) at 1/1000 dilution

Lane 1 : Wild-type HAP1 whole cell lysate

Lane 2 : PRKAG1 (AMPK gamma 1) knockout HAP1 whole cell lysate


Lane 3 : HeLa whole cell lysate

Lysates/proteins at 20 µg per lane.

Predicted band size: 38 kDa

Lanes 1 - 3: Merged signal (red and green). Green - ab32382 observed at 38 kDa. Red - loading control, Mouse anti Vinculin, observed at 125 kDa.

ab32382 was shown to recognize AMPK gamma 1 in wild-type HAP1 cells as signal was lost at the expected MW in PRKAG1 (AMPK gamma 1) knockout cells. Additional cross-reactive bands were observed in the wild-type and knockout cells. Wild-type and PRKAG1 (AMPK gamma 1) knockout samples were subjected to SDS-PAGE. Ab32382 and Mouse anti-Vinculin ([ab9484](#) loading control) were incubated overnight at 4°C at 1/1000 dilution and 1/20000 dilution respectively. Blots were developed with Goat anti-Rabbit IgG H&L (IRDye® 800CW) preabsorbed [ab216773](#) and Goat anti-Mouse IgG H&L (IRDye® 680RD) preabsorbed [ab216776](#) secondary antibodies at 1/20000 dilution for 1 hour at room temperature before imaging.



Western blot - Anti-AMPK gamma 1 antibody [Y307] (ab32382)


All lanes : Anti-AMPK gamma 1 antibody [Y307] (ab32382) at 1/2000 dilution

Lane 1 : Jurkat cell lysate

Lane 2 : HeLa cell lysate

Predicted band size: 38 kDa


Observed band size: 37 kDa



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-AMPK gamma 1 antibody [Y307] (ab32382)

This image shows paraffin embedded human skeletal muscle stained with 1/250 **ab23282**.

Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.



Western blot - Anti-AMPK gamma 1 antibody [Y307] (ab32382)

This image is courtesy of an anonymous Abreview

All lanes : Anti-AMPK gamma 1 antibody [Y307] (ab32382) at 1/2000 dilution

Lane 1 : HeLa whole cell lysate - Control Si

Lane 2 : HeLa whole cell lysate - treated with AMPK gamma 1 siRNA A for 36 hours




Lane 3 : HeLa whole cell lysate - treated with AMPK gamma 1 siRNA B for 36 hours

Secondary

All lanes : HRP-conjugated Donkey polyclonal to rabbit IgG at 1/3000 dilution

Predicted band size: 38 kDa

Why choose a recombinant antibody?

 Research with confidence Consistent and reproducible results	 Long-term and scalable supply Recombinant technology
 Success from the first experiment Confirmed specificity	 Ethical standards compliant Animal-free production

Anti-AMPK gamma 1 antibody [Y307] (ab32382)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors