


Anti-Androgen Receptor antibody [SP107] - N-terminal ab105225

リコンビナント **RabMAb**

★★★★☆ **8 Abreviews** **4 References** 画像数 23

製品の概要

製品名	Anti-Androgen Receptor antibody [SP107] - N-terminal
製品の詳細	Rabbit monoclonal [SP107] to Androgen Receptor - N-terminal
由来種	Rabbit
アプリケーション	適用あり: IHC-P, WB
種交差性	交差種: Mouse, Rat, Human 交差が予測される動物種: Dog, Pig 
免疫原	Synthetic peptide within Human Androgen Receptor aa 300-400 (N terminal). The exact sequence is proprietary. Database link: P10275
ポジティブ・コントロール	IHC-P: Human prostate carcinoma, prostate carcinoma, breast ductal carcinoma, skin squamous cell carcinoma, uterus, endometrial carcinoma, breast, cervix, placenta, prostate, testis, bladder transitional cell carcinoma, hepatocellular carcinoma, lung carcinoma, lung squamous cell carcinoma, pancreatic carcinoma, ovarian carcinoma, Mouse testis, and Rat testis tissue. WB: Rat and Mouse testis tissue lysates; LnCAP and DU 145 cell lysates
特記事項	This product has switched from a hybridoma to recombinant production method on 23rd May 2023. This product is a recombinant monoclonal antibody, which offers several advantages including: - High batch-to-batch consistency and reproducibility - Improved sensitivity and specificity - Long-term security of supply - Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.

バッファー	pH: 7.60 Preservative: 0.1% Sodium azide Constituents: PBS, 1% BSA
精製度	Protein A/G purified
特記事項 (精製)	Purified from TCS by protein A/G.
ポリ/モノ	モノクローナル
クローン名	SP107
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab105225の使用に適用されます**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
IHC-P	★★★★★ (2)	1/400. Boil tissue section in 10mM citrate buffer, pH 6.0 for 10 min followed by cooling at RT for 20 min.
WB	★★★☆☆ (2)	Use a concentration of 1.55 µg/ml. Predicted molecular weight: 99 kDa.

ターゲット情報

機能	Steroid hormone receptors are ligand-activated transcription factors that regulate eukaryotic gene expression and affect cellular proliferation and differentiation in target tissues. Transcription factor activity is modulated by bound coactivator and corepressor proteins. Transcription activation is down-regulated by NR0B2. Activated, but not phosphorylated, by HIPK3 and ZIPK/DAPK3. Isoform 3 and isoform 4 lack the C-terminal ligand-binding domain and may therefore constitutively activate the transcription of a specific set of genes independently of steroid hormones.
組織特異性	Isoform 2 is mainly expressed in heart and skeletal muscle (PubMed:15634333). Isoform 3 is expressed by basal and stromal cells of prostate (at protein level) (PubMed:19244107).
関連疾患	Androgen insensitivity syndrome Spinal and bulbar muscular atrophy X-linked 1 Defects in AR may play a role in metastatic prostate cancer. The mutated receptor stimulates prostate growth and metastases development despite of androgen ablation. This treatment can reduce primary and metastatic lesions probably by inducing apoptosis of tumor cells when they express the wild-type receptor. Androgen insensitivity, partial
配列類似性	Belongs to the nuclear hormone receptor family. NR3 subfamily. Contains 1 nuclear receptor DNA-binding domain.
ドメイン	Composed of three domains: a modulating N-terminal domain, a DNA-binding domain and a C-terminal ligand-binding domain. In the presence of bound steroid the ligand-binding domain interacts with the N-terminal modulating domain, and thereby activates AR transcription factor activity. Agonist binding is required for dimerization and binding to target DNA. The transcription

factor activity of the complex formed by ligand-activated AR and DNA is modulated by interactions with coactivator and corepressor proteins. Interaction with RANBP9 is mediated by both the N-terminal domain and the DNA-binding domain. Interaction with EFCAB6/DJBP is mediated by the DNA-binding domain.

翻訳後修飾

Sumoylated on Lys-388 (major) and Lys-521. Ubiquitinated. Deubiquitinated by USP26. 'Lys-6' and 'Lys-27'-linked polyubiquitination by RNF6 modulates AR transcriptional activity and specificity.

Phosphorylated in prostate cancer cells in response to several growth factors including EGF. Phosphorylation is induced by c-Src kinase (CSK). Tyr-535 is one of the major phosphorylation sites and an increase in phosphorylation and Src kinase activity is associated with prostate cancer progression. Phosphorylation by TNK2 enhances the DNA-binding and transcriptional activity and may be responsible for androgen-independent progression of prostate cancer. Phosphorylation at Ser-83 by CDK9 regulates AR promoter selectivity and cell growth.

Phosphorylation by PAK6 leads to AR-mediated transcription inhibition.

Palmitoylated by ZDHHC7 and ZDHHC21. Palmitoylation is required for plasma membrane targeting and for rapid intracellular signaling via ERK and AKT kinases and cAMP generation.


細胞内局在

Nucleus. Cytoplasm. Predominantly cytoplasmic in unligated form but translocates to the nucleus upon ligand-binding. Can also translocate to the nucleus in unligated form in the presence of RACK1.

製品の状態

There are 2 isoforms produced by alternative splicing. Isoform 1 is also known as: AR-B; isoform 2 is known as AR-A or variant AR45.

画像



All lanes : Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225) at 1/1000 dilution

Lane 1 : Mouse liver lysate

Lane 2 : Rat liver lysate

Lane 3 : Mouse testis lysate

Lane 4 : Rat testis lysate

Lysates/proteins at 20 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/20000 dilution


Predicted band size: 99 kDa

Observed band size: 110 kDa

Exposure time: 23 seconds

Blocking/Diluting buffer: 5% NFDM/TBST

Loading Control: Rabbit monoclonal [EPR16891] to GAPDH
([ab181602](#))



Western blot - Anti-Androgen Receptor antibody
[SP107] - N-terminal ([ab105225](#))

All lanes : Anti-Androgen Receptor antibody [SP107] - N-terminal
([ab105225](#)) at 1.55 µg

Lane 1 : LnCAP (Human prostate carcinoma epithelial cell) cell
lysate

Lane 2 : DU 145 (Human prostate carcinoma epithelial cell) cell
lysate

Lane 3 : Mouse testis tissue lysate

Lysates/proteins at 20 µg per lane.


Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 0.05
µg/ml

Predicted band size: 99 kDa


Observed band size: 120 kDa

Blocking/Diluting buffer and concentration: 5% NFDM /TBST




Immunohistochemistry (Formalin/PFA-fixed paraffin-
embedded sections) - Anti-Androgen Receptor
antibody [SP107] - N-terminal ([ab105225](#))

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Rat testis tissue sections labeling Androgen Receptor with Purified [ab105225](#) at 1/400 dilution (0.37 µg/ml). Heat mediated antigen retrieval was performed Heat mediated antigen retrieval with sodium citrate buffer (pH 6.0, epitope retrieval solution 1) for 30mins. Rabbit specific IHC polymer detection kit HRP/DAB ([ab209101](#)) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Mouse testis tissue sections labeling Androgen Receptor with Purified ab105225 at 1/400 dilution (0.37 µg/ml). Heat mediated antigen retrieval was performed Heat mediated antigen retrieval with sodium citrate buffer (pH 6.0, epitope retrieval solution 1) for 30mins. Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) analysis of Human prostate cancer tissue sections labeling Androgen Receptor with Purified ab105225 at 1/400 dilution (0.37 µg/ml). Heat mediated antigen retrieval was performed Heat mediated antigen retrieval with sodium citrate buffer (pH 6.0, epitope retrieval solution 1) for 30mins. Rabbit specific IHC polymer detection kit HRP/DAB (**ab209101**) was used as the secondary antibody. Negative control: PBS instead of the primary antibody. Hematoxylin was used as a counterstain.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)

Formalin-fixed, paraffin-embedded human prostate carcinoma tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)

Formalin-fixed, paraffin-embedded human prostate adenocarcinoma tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)

Formalin-fixed, paraffin-embedded human breast ductal carcinoma tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)

Formalin-fixed, paraffin-embedded human skin squamous cell carcinoma tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)

Formalin-fixed, paraffin-embedded human uterus tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)

Formalin-fixed, paraffin-embedded human endometrial adenocarcinoma tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.




Formalin-fixed, paraffin-embedded human breast tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)




Formalin-fixed, paraffin-embedded human cervix tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)




Formalin-fixed, paraffin-embedded human placenta tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)




Formalin-fixed, paraffin-embedded human prostate tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)




Formalin-fixed, paraffin-embedded human testis tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)




Formalin-fixed, paraffin-embedded human bladder transitional cell carcinoma tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)

Formalin-fixed, paraffin-embedded human hepatocellular carcinoma tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)

Formalin-fixed, paraffin-embedded human lung adenocarcinoma tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.




Formalin-fixed, paraffin-embedded human lung squamous cell carcinoma tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)



Formalin-fixed, paraffin-embedded human pancreatic adenocarcinoma tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)



Formalin-fixed, paraffin-embedded human ovarian adenocarcinoma tissue stained for Androgen Receptor with ab105225 at 1/400 dilution in immunohistochemical analysis.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Androgen Receptor antibody [SP107] - N-terminal (ab105225)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours

- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors