

Anti-Angiotensinogen antibody [EPR2931] ab108334

リコンビナント RabMAb

★★★★☆ **4 Abreviews** **10 References** 画像数 5

製品の概要

製品名	Anti-Angiotensinogen antibody [EPR2931]
製品の詳細	Rabbit monoclonal [EPR2931] to Angiotensinogen
由来種	Rabbit
特異性	ab108334 is raised against the Human Angiotensinogen protein. This has little homology with mouse or rat Angiotensinogen and in our hands we do not get good results when testing in these species (in western blot the band is weak or absent and at a lower molecular weight than in human samples) therefore we cannot guarantee this product for use in these species. When using this antibody for western blot we recommend blocking with 5% BSA for best results. In our hands, blocking with milk does not give optimal results in WB and some lower molecular weight binding can be observed.
アプリケーション	適用あり: WB, IP, ICC/IF 適用なし: Flow Cyt or IHC-P
種交差性	交差種: Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	Human fetal artery, fetal heart, fetal liver, adult kidney, HepG2, 293T, and plasma lysates; HepG2 cells.
特記事項	This product is a recombinant monoclonal antibody, which offers several advantages including: <ul style="list-style-type: none">- High batch-to-batch consistency and reproducibility- Improved sensitivity and specificity- Long-term security of supply- Animal-free production For more information see here . Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C. Stable for 12 months at -20°C.

バッファー	pH: 7.20 Preservative: 0.01% Sodium azide Constituents: 9% PBS, 40% Glycerol (glycerin, glycerine), 0.05% BSA, 50% Tissue culture supernatant
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR2931
アイソタイプ	IgG

アプリケーション

The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab108334の使用に適用されず**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB	★★★★☆ (1)	1/1000 - 1/10000. Predicted molecular weight: 53 kDa. Block with 5% BSA
IP		1/10 - 1/100.
ICC/IF	★★★★☆ (1)	1/100 - 1/250.

追加情報 Is unsuitable for Flow Cyt or IHC-P.

ターゲット情報

機能 Essential component of the renin-angiotensin system (RAS), a potent regulator of blood pressure, body fluid and electrolyte homeostasis. In response to lowered blood pressure, the enzyme renin cleaves angiotensinogen to produce angiotensin-1 (angiotensin 1-10). Angiotensin-1 is a substrate of ACE (angiotensin converting enzyme) that removes a dipeptide to yield the physiologically active peptide angiotensin-2 (angiotensin 1-8). Angiotensin-1 and angiotensin-2 can be further processed to generate angiotensin-3 (angiotensin 2-8), angiotensin-4 (angiotensin 3-8). Angiotensin 1-7 is cleaved from angiotensin-2 by ACE2 or from angiotensin-1 by MME (neprilysin). Angiotensin 1-9 is cleaved from angiotensin-1 by ACE2.
Angiotensin-2 acts directly on vascular smooth muscle as a potent vasoconstrictor, affects cardiac contractility and heart rate through its action on the sympathetic nervous system, and alters renal sodium and water absorption through its ability to stimulate the zona glomerulosa cells of the adrenal cortex to synthesize and secrete aldosterone.
Angiotensin-3 stimulates aldosterone release.
Angiotensin 1-7 is a ligand for the G-protein coupled receptor MAS1 (By similarity). Has vasodilator and antidiuretic effects (By similarity). Has an antithrombotic effect that involves MAS1-mediated release of nitric oxide from platelets.

組織特異性 Expressed by the liver and secreted in plasma.

関連疾患 Genetic variations in AGT are a cause of susceptibility to essential hypertension (EHT) [MIM:145500]. Essential hypertension is a condition in which blood pressure is consistently higher than normal with no identifiable cause.
Defects in AGT are a cause of renal tubular dysgenesis (RTD) [MIM:267430]. RTD is an

autosomal recessive severe disorder of renal tubular development characterized by persistent fetal anuria and perinatal death, probably due to pulmonary hypoplasia from early-onset oligohydramnios (the Potter phenotype).

配列類似性

Belongs to the serpin family.


翻訳後修飾

Beta-decarboxylation of Asp-34 in angiotensin-2, by mononuclear leukocytes produces alanine. The resulting peptide form, angiotensin-A, has the same affinity for the AT1 receptor as angiotensin-2, but a higher affinity for the AT2 receptor.

細胞内局在

Secreted.

画像



All lanes : Anti-Angiotensinogen antibody [EPR2931] (ab108334) at 1/1000 dilution

Lane 1 : Mouse kidney whole tissue lysate

Lane 2 : Rat kidney whole tissue lysate

Lane 3 : Human kidney whole tissue lysate

Lane 4 : Mouse liver whole tissue lysate

Lane 5 : Rat liver whole tissue lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat anti-rabbit IgG, (H+L), peroxidase conjugated at 1/2000 dilution

Predicted band size: 53 kDa

Exposure time: 1 minute

Ab108334 targeting angiotensinogen in various mouse, rat and human tissues (see lane descriptions).

Blocking buffer for the first lanes 1-5: 5% NFD/MTBST; the second lanes 1-5: BSA/TBST.



All lanes : Anti-Angiotensinogen antibody [EPR2931] (ab108334) at 1/1000 dilution

All lanes : Human fetal liver whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary


All lanes : Goat anti-rabbit IgG (H+L) peroxidase conjugated at 1/2000 dilution

Predicted band size: 53 kDa

Exposure time: 3 minutes

Lane 1 blocking buffer: NFDm/TBST

Lane 2 blocking buffer: BSA/TBST



All lanes : Anti-Angiotensinogen antibody [EPR2931] (ab108334) at 1/1000 dilution

Lane 1 : Human fetal artery lysate

Lane 2 : Human fetal heart lysate

Lane 3 : Human fetal liver lysate


Lane 4 : HepG2 lysate

Lane 5 : 293T lysate

Lane 6 : Human plasma lysate

Lysates/proteins at 10 µg per lane.

Predicted band size: 53 kDa



ab108334, at 1/100 dilution, staining Angiotensinogen in HepG2 cells by Immunofluorescence.

Immunocytochemistry/ Immunofluorescence - Anti-Angiotensinogen antibody [EPR2931] (ab108334)

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-Angiotensinogen antibody [EPR2931]
(ab108334)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors