




Anti-Aspartate Aminotransferase antibody ab85857

3 References [画像数 3](#)

製品の概要

製品名	Anti-Aspartate Aminotransferase antibody
製品の詳細	Goat polyclonal to Aspartate Aminotransferase
由来種	Goat
アプリケーション	適用あり: WB, IHC-P
種交差性	交差種: Human 交差が予測される動物種: Rat, Pig, Chimpanzee, Cynomolgus monkey, Orangutan 
免疫原	Synthetic peptide: RSYRYWDAEKR by a Cysteine residue linker, corresponding to C terminal amino acids 157-167 of Human Aspartate Aminotransferase <div>  Run BLAST with  Run BLAST with </div>
ポジティブ・コントロール	Human kidney and heart tissue, Human Liver lysate.
特記事項	<p>The Life Science industry has been in the grips of a reproducibility crisis for a number of years. Abcam is leading the way in addressing this with our range of recombinant monoclonal antibodies and knockout edited cell lines for gold-standard validation. Please check that this product meets your needs before purchasing.</p> <p>If you have any questions, special requirements or concerns, please send us an inquiry and/or contact our Support team ahead of purchase. Recommended alternatives for this product can be found below, along with publications, customer reviews and Q&As</p>

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Upon delivery aliquot and store at -20°C. Avoid freeze / thaw cycles.
バッファー	pH: 7.30 Preservative: 0.02% Sodium azide Constituents: Tris buffered saline, 0.5% BSA
精製度	Immunogen affinity purified
ポリ/モノ	ポリクローナル
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee Abpromise保証は、 次のテスト済みアプリケーションにおけるab85857の使用に適用されます
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
WB		Use a concentration of 0.01 - 0.1 µg/ml. Predicted molecular weight: 46 kDa.
IHC-P		Use a concentration of 2.5 - 5 µg/ml.

ターゲット情報


機能	Plays a key role in amino acid metabolism.
配列類似性	Belongs to the class-I pyridoxal-phosphate-dependent aminotransferase family.
細胞内局在	Cytoplasm.

画像



ab85857 at 5 µg/ml staining Aspartate Aminotransferase in formalin fixed and paraffin embedded Human Kidney tissue.

Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Aspartate Aminotransferase antibody (ab85857)



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-Aspartate Aminotransferase antibody (ab85857)

ab85857 at 5 µg/ml staining Aspartate Aminotransferase in formalin fixed and paraffin embedded Human heart tissue.



Western blot - Anti-Aspartate Aminotransferase antibody (ab85857)

Anti-Aspartate Aminotransferase antibody (ab85857) at 0.01 µg/ml + Human Liver lysate at 35 µg

Developed using the ECL technique.

Predicted band size: 46 kDa

Observed band size: 46 kDa

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors