

Anti-ATG16L1 (phospho S278) antibody [EPR19016] ab195242

リコンビナント RabMAb[®]

★★★★★ **5 Abreviews** **9 References** [画像数 9](#)

製品の概要

製品名	Anti-ATG16L1 (phospho S278) antibody [EPR19016]
製品の詳細	Rabbit monoclonal [EPR19016] to ATG16L1 (phospho S278)
由来種	Rabbit
アプリケーション	適用あり: ICC/IF, IHC-P, Dot blot, WB
種交差性	交差種: Mouse, Human
免疫原	Synthetic peptide. This information is proprietary to Abcam and/or its suppliers.
ポジティブ・コントロール	WB: HEK-293 overexpressing ATG16L1 (WT) whole cell lysate. IHC-P: Human muscle tissue.
特記事項	Co-immunization performed with both peptides, clone obtained by screening with peptide 1. Our RabMAb [®] technology is a patented hybridoma-based technology for making rabbit monoclonal antibodies. For details on our patents, please refer to RabMAb[®] patents .

製品の特性

製品の状態	Liquid
保存方法	Shipped at 4°C. Store at +4°C short term (1-2 weeks). Upon delivery aliquot. Store at -20°C long term. Avoid freeze / thaw cycle.
バッファー	pH: 7.2 Preservative: 0.01% Sodium azide Constituents: 0.05% BSA, 40% Glycerol, PBS
精製度	Protein A purified
ポリ/モノ	モノクローナル
クローン名	EPR19016
アイソタイプ	IgG

アプリケーション


The Abpromise guarantee **Abpromise保証は、次のテスト済みアプリケーションにおけるab195242の使用に適用されず**
アプリケーションノートには、推奨の開始希釈率がありますが、適切な希釈率につきましてはご検討ください。

アプリケーション	Abreviews	特記事項
ICC/IF		1/150.
IHC-P		Use a concentration of 3 µg/ml. Perform heat mediated antigen retrieval with citrate buffer pH 6 before commencing with IHC staining protocol.
Dot blot		1/1000.
WB	★★★★★ (5)	Use at an assay dependent concentration. Predicted molecular weight: 68 kDa. For optimal WB signal, we recommend using 10X Blocking Buffer (ab126587).

ターゲット情報

機能	Plays an essential role in autophagy: interacts with ATG12-ATG5 to mediate the conjugation of phosphatidylethanolamine (PE) to LC3 (MAP1LC3A, MAP1LC3B or MAP1LC3C), to produce a membrane-bound activated form of LC3 named LC3-II. Thereby, controls the elongation of the nascent autophagosomal membrane.
関連疾患	Inflammatory bowel disease 10
配列類似性	Belongs to the WD repeat ATG16 family. Contains 7 WD repeats.
翻訳後修飾	Proteolytic cleavage by activated CASP3 leads to degradation and may regulate autophagy upon cellular stress and apoptotic stimuli.
細胞内局在	Cytoplasm. Preautophagosomal structure membrane. Recruited to omegasomes membranes by WIPI2. Omegasomes are endoplasmic reticulum connected structures at the origin of preautophagosomal structures. Localized to preautophagosomal structure (PAS) where it is involved in the membrane targeting of ATG5. Localizes also to discrete punctae along the ciliary axoneme.
製品の状態	There are 4 isoforms produced by alternative splicing.

画像



Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ATG16L1 (phospho S278) antibody [EPR19016] (ab195242)

IHC images of vessel staining of ab195242, ATG16L1 (phospho S278), in sections of formalin-fixed paraffin-embedded normal human skeletal muscle tissue*, performed on a Leica BOND™ system using a modified protocol F.


Both sections were pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6, epitope retrieval solution 1) for 20mins. One section was then treated with 200 enzyme units of alkaline phosphatase (AP+) for 1 hour at 37°C; and the other in buffer containing no alkaline phosphatase (AP-) for 1 hour at 37°C. The sections were then incubated with ab195242, 3µg/ml, for 15 mins at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The sections were then counterstained with haematoxylin and mounted with DPX.

Identical assays were also performed using detection system-only (no primary antibody) as reagent controls (data not shown), to ensure that staining seen was a result of the binding of the primary antibody.

The absence of staining in the AP+ tissue compared to the AP- tissue adds further evidence of phospho specificity for this antibody.

For other IHC staining systems (automated and non-automated) customers should optimize variable parameters such as antigen retrieval conditions, primary antibody concentration and antibody incubation times.

*Tissue obtained from the Human Research Tissue Bank, supported by the NIHR Cambridge Biomedical Research Centre




Immunohistochemistry (Formalin/PFA-fixed paraffin-embedded sections) - Anti-ATG16L1 (phospho S278) antibody [EPR19016] (ab195242)

This image is courtesy of Dr. Ryan Russell (University of Ottawa).

IHC images of mice quadriceps showing either pATG16L1 or LC3B staining

Mice were fed ad libitum or starved for 16 hours. Quadriceps muscle were immediately harvested and fixed in 10% formalin for 2 days. The samples were then paraffin embedded, sectioned into 4µm thick slices, and mounted onto glass microscope slides. Slides were stained with primary antibody overnight at 4°C: LC3B 1/1000, pATG16L1 (ab195242) 1/300. Secondary antibody: Alexa Fluor 555 anti-rabbit, 1/1000.




Immunocytochemistry/ Immunofluorescence - Anti-ATG16L1 (phospho S278) antibody [EPR19016] (ab195242)

This image is courtesy of Dr. Ryan Russell (University of Ottawa).

IF showing pATG16L1 (red) and total ATG16L1 (green):

Polyclonal population of ATG16L1 KO and HA-ATG16L1 reconstituted cells were starved of amino acid for 1 hour and stained. Blocking buffer used for pATG16L1 staining: 0.1% BSA, 1x abcam blocking buffer [ab126587](#), diluted in PBS. Anti-pATG16L1 (ab195242) concentration: 1/150. Anti-ATG16L1, concentration: 1/200 Secondary antibody (Alexa Fluor 647/488) concentration: 1/1000.




Immunocytochemistry/ Immunofluorescence - Anti-ATG16L1 (phospho S278) antibody [EPR19016] (ab195242)

This image is courtesy of Dr. Ryan Russell (University of Ottawa).

IF showing pATG16L1 (red), LC3B (green) and p62 (white):


MEF cells were amino acid starved for 1 hour. Blocking buffer used for pATG16L1 (ab195242): 0.1% BSA, 1x abcam blocking buffer (**ab126587**), diluted in PBS. Anti-pATG16L1 (ab195242) concentration: 1/150. Secondary antibody (Alexa Fluor 647) concentration: 1/1000



Western blot - Anti-ATG16L1 (phospho S278) antibody [EPR19016] (ab195242)

This image is courtesy of Dr Ryan Russell (University of Ottawa).

HCT116 wild-type and ATG16L1 knockout cells were incubated with either complete media or amino acid deficient DMEM for 3 hours. 5ug of whole cell lysate were resolved by SDS-PAGE on a 6%-18% gradient gel, then transferred onto PVDF membrane. Membrane was blocked in 10X blocking buffer (Cat # **ab126587**) diluted in TBS solution for 30 minutes; incubated with 1:1000 primary antibody in 2.5% BSA TBST solution overnight at 4°C ; incubated with 1:15000 secondary antibody in 2% milk TBST solution for 45 minutes. Immobilon ECL was applied for 1 minute then imaged with film.



Western blot - Anti-ATG16L1 (phospho S278) antibody [EPR19016] (ab195242)

All lanes : Anti-ATG16L1 (phospho S278) antibody [EPR19016] (ab195242) at 1/1000 dilution

Lane 1 : HEK-293 (human epithelial cell line from embryonic kidney) transfected with an empty vector (vector control), containing a myc-His-tag®, whole cell lysate

Lane 2 : HEK-293 (human epithelial cell line from embryonic kidney) transfected with ATG16L1 (WT) expression vector containing a myc-His-tag®, whole cell lysate

Lane 3 : HEK-293 (human epithelial cell line from embryonic kidney) transfected with ATG16L1 (WT) expression vector containing a myc-His-tag®, followed by treatment with alkaline phosphatase for 1 hour, whole cell lysate

Lane 4 : HEK-293 (human epithelial cell line from embryonic kidney) transfected with ATG16L1 S278A expression vector containing a myc-His-tag®, whole cell lysate

Lysates/proteins at 10 µg per lane.

Secondary

All lanes : Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100000 dilution


Developed using the ECL technique.

Predicted band size: 68 kDa

Observed band size: 68 kDa

Exposure time: 3 minutes


Blocking/Dilution buffer: 5% NFDM/TBST.



Western blot - Anti-ATG16L1 (phospho S278) antibody [EPR19016] (ab195242)

This image is courtesy of Dr Ryan Russell (University of Ottawa).

Cells were treated with Torin-1 (200uM) for 1 hour. Membrane was blocked in 10X blocking buffer (Cat # [ab126587](#)) diluted in PBS solution for 30 minutes; incubated with 1/1000 primary antibody in 2.5% BSA TBST solution overnight at 4°C ; incubated with 1/7000 secondary antibody in 2% milk TBST solution for 1 hour.



Dot Blot - Anti-ATG16L1 (phospho S278) antibody [EPR19016] (ab195242)

Dot blot analysis of ATG16L1 (phospho S278) labeled with ab195242 at 1/1,000 dilution.

Lane 1: Mouse ATG16L1 (phospho S278) peptide;

Lane 2: Mouse ATG16L1 non-phospho peptide;

Goat Anti-Rabbit IgG H&L (HRP) ([ab97051](#)) at 1/100,000 dilution was used as secondary antibody.

Blocking/Dilution buffer: 5% NFDM/TBST.

Exposure time: 3 minutes.

Why choose a recombinant antibody?



Research with confidence
Consistent and reproducible results



Long-term and scalable supply
Recombinant technology



Success from the first experiment
Confirmed specificity



Ethical standards compliant
Animal-free production

Anti-ATG16L1 (phospho S278) antibody [EPR19016]
(ab195242)

Please note: All products are "FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES"

Our Abpromise to you: Quality guaranteed and expert technical support

- Replacement or refund for products not performing as stated on the datasheet
- Valid for 12 months from date of delivery
- Response to your inquiry within 24 hours
- We provide support in Chinese, English, French, German, Japanese and Spanish
- Extensive multi-media technical resources to help you
- We investigate all quality concerns to ensure our products perform to the highest standards

If the product does not perform as described on this datasheet, we will offer a refund or replacement. For full details of the Abpromise, please visit <https://www.abcam.co.jp/abpromise> or contact our technical team.

Terms and conditions

- Guarantee only valid for products bought direct from Abcam or one of our authorized distributors